



NPCA WATER QUALITY MONITORING PROGRAM:
SUMMARY REPORT OF THE YEAR 2022



SEPTEMBER 2023

TABLE OF CONTENTS

<u>1.0 INTRODUCTION</u>	6
<u>2.0 SURFACE WATER QUALITY MONITORING PROGRAM</u>	6
2.1 CHEMICAL MONITORING	6
2.1.1 NIAGARA RIVER AOC TRIBUTARY MONITORING PROGRAM	6
2.1.2 PROVINCIAL WATER QUALITY MONITORING NETWORK	6
2.1.3 OTHER WATER QUALITY MONITORING PROGRAMS	7
2.2 BIOLOGICAL MONITORING	7
<u>3.0 SURFACE WATER QUALITY INDICATOR PARAMETERS</u>	8
3.1 CHLORIDE	8
3.2 NITRATE	8
3.3 TOTAL PHOSPHORUS	9
3.4 SUSPENDED SOLIDS	9
3.5 COPPER	9
3.6 LEAD	9
3.7 ZINC	9
3.8 ESCHERICHIA COLI	10
3.9 BIOLOGICAL ASSESSMENTS BENTHIC INVERTEBRATES	10
<u>4.0 SURFACE WATER QUALITY MONITORING RESULTS</u>	10
4.1 TREND ANALYSIS	12
4.2 WELLAND RIVER WATERSHED	13
4.2.1 WELLAND RIVER: WATER QUALITY INDEX	13
4.2.2 WELLAND RIVER: HILSENHOFF BIOTIC INDEX RESULTS	18
4.2.3 WELLAND RIVER: KEY FINDINGS	18
4.3 WELLAND RIVER TRIBUTARIES	19
4.3.1 WELLAND RIVER TRIBUTARIES: WATER QUALITY INDEX	19
4.3.2 WELLAND RIVER TRIBUTARIES: HILSENHOFF BIOTIC INDEX RESULTS	23
4.3.3 WELLAND RIVER TRIBUTARIES: KEY FINDINGS	23
4.4 TWENTY MILE CREEK WATERSHED	24
4.4.1 TWENTY MILE CREEK WATERSHED: WATER QUALITY INDEX	24
4.4.2 TWENTY MILE CREEK WATERSHED: HILSENHOFF BIOTIC INDEX RESULTS	27
4.4.3 TWENTY MILE CREEK: KEY FINDINGS	27
4.5 LAKE ONTARIO TRIBUTARIES	28
4.5.1 LAKE ONTARIO TRIBUTARIES: WATER QUALITY INDEX	28
4.5.2 LAKE ONTARIO TRIBUTARIES: HILSENHOFF BIOTIC INDEX RESULTS	35
4.5.3. LAKE ONTARIO TRIBUTARIES: KEY FINDINGS	35
4.6 NIAGARA RIVER TRIBUTARIES	36
4.6.1 NIAGARA RIVER TRIBUTARIES: WATER QUALITY INDEX	36
4.6.2 NIAGARA RIVER TRIBUTARIES: HILSENHOFF BIOTIC INDEX RESULTS	39
4.6.3 NIAGARA RIVER TRIBUTARIES: KEY FINDINGS	39
4.7 LAKE ERIE TRIBUTARIES	39
4.7.1 LAKE ERIE TRIBUTARIES: WATER QUALITY INDEX	39
4.7.2 LAKE ERIE TRIBUTARIES: HILSENHOFF BIOTIC INDEX RESULTS	43
4.7.3 LAKE ERIE TRIBUTARIES: KEY FINDINGS	43

<u>5.0 GROUNDWATER QUALITY MONITORING PROGRAM</u>	43
5.1 PROVINCIAL GROUNDWATER MONITORING NETWORK	43
5.1.1 GROUNDWATER LEVELS	46
5.1.2 GROUNDWATER CHEMISTRY	46
5.1.3 GROUNDWATER KEY FINDINGS	51
5.2 WATER WELL DECOMMISSIONING PROGRAM	52
<u>6.0 OTHER PROJECTS</u>	53
6.1 HAMILTON INTERNATIONAL AIRPORT	53
6.2 GLANBROOK LANDFILL	53
6.3 UPPER TWELVE MILE CREEK TEMPERATURE STUDY	54
6.4 LAKE NIAPENCO PERFLUORINIATED COMPOUND MONITORING	58
6.5 CANADA ONTARIO AGREEMENT CLIMATE CHANGE MONITORING PROJECT	61
6.6 CONSERVATION AREA WATER QUALITY MONITORING	62
6.7 LOWER TWELVE MILE CREEK TOTAL PCBs MONITORING	66
6.9 NPCA DATA REQUESTS	68
<u>7.0 CONCLUSIONS</u>	69
<u>8.0 RECOMMENDATIONS</u>	70
<u>9.0 REFERENCES CITED</u>	71
<u>10.0 ACKNOWLEDGEMENTS</u>	73

APPENDIX A: SURFACE WATER QUALITY INDEX RATINGS: 2018-2022

APPENDIX B: MEDIAN CONCENTRATIONS OF CHLORIDE FOR NPCA MONITORING STATIONS

APPENDIX C: MEDIAN CONCENTRATIONS OF COPPER FOR NPCA MONITORING STATIONS

APPENDIX D: MEDIAN CONCENTRATIONS OF E. COLI FOR NPCA MONITORING STATIONS

APPENDIX E: MEDIAN CONCENTRATIONS OF LEAD FOR NPCA MONITORING STATIONS

APPENDIX F: MEDIAN CONCENTRATIONS OF NITRATE FOR NPCA MONITORING STATIONS

APPENDIX G: MEDIAN CONCENTRATIONS OF TOTAL PHOSPHORUS FOR NPCA MONITORING STATIONS

APPENDIX H: MEDIAN CONCENTRATIONS OF TOTAL SUSPENDED SOLIDS FOR NPCA MONITORING STATIONS

APPENDIX I: MEDIAN CONCENTRATIONS OF ZINC FOR NPCA MONITORING STATIONS

APPENDIX J: HILSENHOFF BIOTIC INDEX WATER QUALITY ASSESSMENTS 2018-2022

APPENDIX K: GROUNDWATER LEVELS

Table of Acronyms

AOC	Area of Concern
BMPs	Best Management Practices
CCME	Canadian Council of Ministers of the Environment
CWQG	Canadian Water Quality Guidelines
ECCC	Environment and Climate Change Canada
EMRB	Environmental Monitoring and Reporting Branch
HIA	Hamilton International Airport
HBI	Hilsenhoff Biotic Index
MECP	Ministry of Environment Conservation and Parks
NPCA	Niagara Peninsula Conservation Authority
PFC	Perfluorinated Compound
PGMN	Provincial Groundwater Monitoring Network
PWQMN	Provincial Water Quality Monitoring Network
PWQO	Provincial Water Quality Objective
OBBN	Ontario Benthos Biomonitoring Network
ODWS	Ontario Drinking Water Standards
OGS	Ontario Geologic Survey
OPG	Ontario Power Generation
RMN	Regional Municipality of Niagara
WQI	Water Quality Index for CCME

EXECUTIVE SUMMARY

The Niagara Peninsula Conservation Authority's (NPCA) Water Quality Monitoring Program was implemented in 2001 and is operated in partnership with the Ontario Ministry of Environment, Conservation and Parks (MECP), Regional Municipality of Niagara, Haldimand County, and the City of Hamilton. Through these partnerships the NPCA collects water quality samples at 80 surface water stations and 46 groundwater wells located throughout the NPCA watershed. The NPCA utilizes both chemical and biological approaches to evaluate the surface water quality. Surface water quality samples are analysed for several indicators such as chloride, nutrients, *E. coli*, suspended solids, and metals. Surface water results are used to calculate the Canada Council of Ministers of Environment Water Quality Index. This index is a Canada-wide standard for reporting water quality information. The NPCA also evaluates water quality in the watershed by sampling the aquatic animals at most of the NPCA surface water quality stations using the Ontario Biological Benthos Monitoring (OBBN) protocol. The abundance and diversity of animals living in the watercourse provides a biological assessment of the water quality. Groundwater samples are evaluated by comparing monitoring results to the MECP's Drinking Water Standards (ODWS).

For surface water, the monitoring results indicate most of the NPCA's watersheds have poor water quality. Total phosphorus, *E. coli*, suspended solids, and chlorides from non-point sources (agricultural/livestock runoff, faulty septic systems) and point sources (combined sewer overflow, urban stormwater) continue to be the major causes of impairment in the NPCA watershed. Watercourses under the direct influence from the Great Lakes and Niagara River such as the Lower Welland River have higher water quality ratings. Watershed tributaries that are strongly influenced by groundwater discharge like the Upper Twelve Mile Creek Effingham portion and Upper Welland River as well as the tributaries found with substantial natural landscapes like Point Abino Drain and Beaver Creek (Fort Erie) continue to have the best water quality ratings in the NPCA watershed.

The ambient groundwater geochemistry in NPCA jurisdiction was found to have some health and aesthetic ODWS exceedances. Health ODWS exceedances have been investigated by MECP and are attributed to natural conditions of the groundwater but in the case of sodium exceedances, the local Health Units are periodically notifying local physicians of elevated sodium levels in the groundwater for them to advise their patients with sodium-restricted diets. The NPCA recommends that residents using groundwater regularly test their well water and use the appropriate water treatment.

The Water Quality Monitoring Program continues to provide valuable information about the health of the NPCA watershed. Often the way the land is managed is reflected in the health of our water resources. The fact that the water quality is generally poor in the NPCA watershed has been caused by decades of environmental degradation. However, water quality improvement programs that refine how nutrients are managed, increase riparian buffers, and expand forest cover can begin to address these impacts. It will likely take many years of implementing these programs before the water quality in the NPCA watershed improves to the point where it is able to meet federal and provincial water quality guidelines and objectives. As such, it is recommended that the NPCA continue to monitor both our surface water and groundwater to ensure that there is up-to-date current water quality information available, be able to quantify trends, and continue to identify sources of contamination within the NPCA watershed.

NPCA WATER QUALITY MONITORING PROGRAM: SUMMARY REPORT FOR THE YEAR 2022

1.0 INTRODUCTION

The NPCA Water Quality Monitoring Program was initiated in 2001. Before 2001, the NPCA was involved in numerous water quality related initiatives but did not have a dedicated monitoring program. The NPCA has since established an extensive network of monitoring stations located throughout the watershed with the purpose of gathering long-term water quality data for both surface water and groundwater. This network represents the largest and most comprehensive water quality monitoring program in the Niagara Peninsula. The NPCA monitoring network is operated in partnership with the MECP, Regional Municipality of Niagara (RMN), Haldimand County and City of Hamilton. The main objective of the NPCA Water Quality Monitoring Program is to assess water quality in local watersheds using a network of chemical and biological monitoring stations. The purpose of this Annual Report is to summarize the water quality data collected from these monitoring stations and provide recommendations for future monitoring and stewardship initiatives.

2.0 SURFACE WATER QUALITY MONITORING PROGRAM

2.1 CHEMICAL MONITORING

In 2022, the NPCA monitored surface water quality at 80 stations covering 52 watersheds. Grab samples are collected monthly during the ice-free season and analyzed for several parameters including nutrients, metals, bacteria, suspended solids, and general chemistry. The chemical monitoring program is mainly funded through the municipal levy, however the NPCA does receive additional support for lab analysis from the following partners: RMN, MECP, ECCC and the City of Hamilton. These are described in detailed below.

2.1.1 NIAGARA RIVER AOC TRIBUTARY MONITORING PROGRAM

The Niagara River was listed as a Great Lakes' Area of Concern (AOC) in the late 1980s due to habitat loss and contaminants in water. A Remedial Action Plan (RAP) was first developed and implemented beginning in 1993. This monitoring program was established in 2003 out of a recommendation in the 1995 Niagara River Stage 2 RAP Report (MOE and EC 1995) which outlined 37 recommended remedial actions to restore the health of the watershed. Recommendation #29 was to develop and implement a Welland River and Niagara River tributaries monitoring program to monitor rural non-point sources of pollution, establish a baseline of water quality conditions at selected tributaries, and track the effectiveness of stewardship efforts in the watershed (MOE and EC 1995). Stations were selected to both overlap with historic stations and fill data gaps where required. The monitoring program was implemented as part of an agreement with the federal and provincial government until 2016. Given the scope of the AOC was recently focused on the Niagara River proper (as per guidance in the Great Lake Water Quality Agreement), the NPCA began funding the sample collection and lab analysis for these monitoring stations in 2017.

2.1.2 PROVINCIAL WATER QUALITY MONITORING NETWORK

In 2003 a partnership was established with the MECP through the Provincial Water Quality Monitoring Network (PWQMN) whereby NPCA staff collect monthly water samples at stations located within the NPCA watershed and the MECP provides

laboratory services. The PWQMN was established in 1964 to collect surface water quality information from rivers and streams at strategic locations throughout Ontario. Over time, stations have been added and discontinued in response to changing MECP and program-specific needs. The NPCA has 13 PWQMN stations which are located on the Black Creek (Fort Erie), Welland River (West Lincoln & Welland), Twenty Mile Creek (West Lincoln and Lincoln), Forty Mile Creek (Grimsby), Four Mile Creek (Niagara-on-the Lake), and Twelve Mile Creek (Pelham & St. Catharines).

2.1.3 OTHER WATER QUALITY MONITORING PROGRAMS

Several watersheds are monitored through other water quality monitoring programs. In 2002 a monitoring agreement was established with the City of Hamilton whereby NPCA staff collect monthly water samples at eleven stations located within the City of Hamilton's municipal boundaries and the City of Hamilton provides laboratory services. This laboratory partnership was cancelled January 2018 by the City of Hamilton. However, after the cancellation of this program the NPCA continued to fund the lab analysis for these monitoring stations. The NPCA is also involved in monitoring at the Hamilton International Airport and the Glanbrook Landfill. For further details, please refer to section 6.0. In 2003 a similar monitoring arrangement was established with the RMN whereby NPCA staff collect water samples at ten stations located within the Niagara Regional Municipal boundary of the NPCA watershed and the RMN provides laboratory services.

2.2 BIOLOGICAL MONITORING

The NPCA also monitors surface water quality using benthic invertebrates as indicators of stream health. Water quality monitoring has historically relied heavily upon chemical testing as a means of measuring the quality of water but the advantages of biological monitoring using benthic invertebrates as indicators of water quality are well documented (Griffiths 1999, Jones *et al.* 2005). Due to their restricted mobility and habitat preferences benthic invertebrates usually remain in a localized area. As a result, they are continuously subjected to the effects of all pollutants and environmental stream conditions over time, and as such can provide a broad overview of water quality related problems. They are abundant in all types of aquatic systems and can be easily collected and identified.

The NPCA has been using benthic invertebrates as indicators of water quality since 1995 and is a leader in the field of biological monitoring in the Niagara Peninsula. Benthic invertebrate samples are collected annually during the spring and fall seasons using the Ontario Benthos Biomonitoring Network (OBBN) protocol. The OBBN provides a standardized benthic invertebrate sampling protocol for the province of Ontario and provides a biological complement to the chemistry based PWQMN. The NPCA is an active participant in the development of the OBBN and is providing on-going research support in the upper Twelve Mile Creek watershed. The NPCA uses the Hilsenhoff Biotic Index (HBI) to summarize biological monitoring data in this report.

3.0 SURFACE WATER QUALITY INDICATOR PARAMETERS

The indicator parameters described in the following sections best reflect the range of water quality issues that are likely encountered in the NPCA watershed and are most useful in assessing relative stream health. These indicator parameters and their respective surface water quality objectives are summarized in **Table 1**.

Table 1: Summary of surface water quality indicator parameters

INDICATOR PARAMETER	OBJECTIVE	REFERENCE
Chloride	120 mg/L (Chronic)	CWQG (CCME 2011)
Nitrate	2.93 mg/L	CWQG (CCME 2003)
Total phosphorus	30 µg/L	PWQO (MOE 1994)
Suspended solids	35 mg/L	CWQG (CCME 2002)
Copper	5 µg/L	PWQO (MOE 1994)
Lead	5 µg/L	PWQO (MOE 1994)
Zinc	20 µg/L	PWQO (MOE 1994)
<i>Escherichia coli</i>	100 counts/100 mL	PWQO (MOE 1994)
Benthic invertebrates	Unimpaired	HBI (Hilsenhoff 1987)

3.1 CHLORIDE

Chloride is a naturally occurring substance found in all waters. Chloride can be toxic to aquatic organisms with acute toxic effects at high concentrations and chronic effects on growth and reproduction at lower concentrations. Chloride ions are conservative, which means that they are not degraded in the aquatic environment and tend to remain in solution. Chloride is extensively used in the form of sodium chloride and calcium chloride for salting of roadways and ice removal during the winter season. Other anthropogenic or human-derived sources of chloride include sewage, animal waste, storm and irrigation drainage, fertilizers, and industrial effluent. The Canadian Water Quality Guidelines (CWQG) for the Protection of Aquatic Life recommend that long-term or chronic chloride concentrations should not exceed 120 mg/L in surface water (CCME 2011).

3.2 NITRATE

Nitrate is the most common form of nitrogen that occurs in surface water. In aerobic or oxygen-rich water, bacteria convert ammonium and nitrite to nitrate through a process known as nitrification. In anaerobic or oxygen-depleted water, the process is reversed through denitrification. The nitrate ion is the most stable form of nitrogen in water and does not tend to combine with other ions in solution. Nitrate can be toxic to aquatic organisms and elevated concentrations contribute to excessive plant and algae growth in surface water. Anthropogenic sources of nitrate include sewage discharges, animal waste, fertilizers, and pesticides. The CWQG for the Protection of Aquatic Life recommend that nitrate-nitrogen concentrations should not exceed 2.93 mg/L in surface water (CCME 2003).

3.3 TOTAL PHOSPHORUS

Phosphorus is a natural element found in rocks, soils and organic material and is an essential nutrient for plant growth. Phosphorus clings tightly to soil particles and is often associated with suspended sediment. Excessive phosphorus concentrations stimulate the overgrowth and decomposition of plants and algae. The decomposition of organic matter in turn depletes dissolved oxygen concentrations and stresses aquatic organisms such as fish and benthic invertebrates. Total phosphorus is a measure of all forms of phosphorus in a water sample and includes biologically accessible phosphates. Anthropogenic sources of phosphorus include fertilizers, pesticides, and sewage discharges. The interim Ontario Provincial Water Quality Objective (PWQO) for total phosphorus in streams and rivers is 30 µg/L (MOE 1994).

3.4 TOTAL SUSPENDED SOLIDS

Total suspended solids (TSS) are a measure of undissolved solid material in surface water and usually consist of silt, clay, plankton, and fine particles of organic and inorganic matter. Sources of suspended solids include soil erosion, stormwater, wastewater, and industrial effluent. Fine particles are significant carriers of phosphorus, metals and other contaminants. Concentrations of suspended solids vary seasonally and often peak during rain events. Due to natural variability in surface water there is currently no water quality guideline for suspended solids in Ontario. High concentrations of suspended solids in surface water can negatively impact aquatic organisms. The Canadian Water Quality Guidelines provides a narrative guideline for TSS: the maximum increase of TSS should be no more than 25 mg/L from background concentrations (with NPCA using a background TSS concentration of 10 mg/L determined using data from the NPCA jurisdiction; CCME 2002). Therefore, a concentration of 35 mg/L was used as the NPCA's guideline.

3.5 COPPER

Copper is an essential trace element that is toxic to aquatic organisms at elevated concentrations. In surface water copper tends to bind with organic matter and accumulate in streambed sediment. Natural sources are wind-blown dust, decaying vegetation and from forest fires. Anthropogenic sources of copper include industrial wastewater, sewage discharges and pesticides. The interim PWQO for copper is 5 µg/L based on >20 mg/L Hardness as CaCO₃ (MOE 1994).

3.6 LEAD

Lead is a non-essential trace element that is toxic to aquatic organisms at elevated concentrations. Lead tends to bioaccumulate and can affect the central nervous system. Lead occurs naturally in the environment. However, most lead concentrations that are found in the environment are a result of human activities. Anthropogenic sources of lead include industrial wastewater, sewage discharges, municipal waste incineration, fertilizers, and pesticides. The interim PWQO for lead is 5 µg/L based on >80 mg/L Hardness as CaCO₃ (MOE 1994).

3.7 ZINC

Zinc is an essential trace element that is toxic to aquatic organisms at elevated concentrations. In surface water zinc tends to bind with organic matter and accumulate in streambed sediment. Zinc occurs naturally in air, water, and soil. Anthropogenic

sources of zinc include industrial wastewater, sewage discharges and stormwater runoff. The interim PWQO for zinc is 20 µg/L (MOE 1994).

3.8 *ESCHERICHIA COLI*

Escherichia coli (*E. coli*) is a type of fecal coliform bacteria that is commonly found in the intestines of warm-blooded animals and humans. *E. coli* is used as an indicator for the presence of sewage or animal waste in surface water, and the possible presence of pathogens (Tchobanoglous & Schroeder 1987). The PWQO for *E. coli* is 100 counts per 100 mL (MOE 1994).

3.9 BIOLOGICAL ASSESSMENTS: BENTHIC INVERTEBRATES

Benthic invertebrates are the larger organisms inhabiting the substrate of watercourses for at least part of their life cycle. As a rule, benthic invertebrates include those species whose body width exceeds 500 microns. Examples of benthic invertebrate species that are commonly found in the NPCA watershed include clams, snails, leeches, worms, and the larval stages of dragonflies, stoneflies, caddisflies, mayflies, and beetles.

The NPCA collects benthic samples during the spring and fall seasons each year at approximately 30 monitoring sites. Once collected, counted, and preserved, the benthic invertebrates are identified to family level and various statistics were calculated. For this 2020 Report the Hilsenhoff Biotic Index (HBI) was calculated for each sample site. The HBI estimates the overall tolerance of the community in a sampled area, weighted by the relative abundance of each family taxonomic group. Organisms are assigned a tolerance value based on those provided Hilsenhoff (1987). Water quality is classified as gradient from *excellent* to *very poor* in to recognize the occurrence of organisms whose environmental requirements and tolerances match those which would be expected at the site without the input of environmental stresses to those with the organisms found are less sensitive. Therefore, more tolerant to environmental stresses than organisms which would have historically occurred. The benthic population at an impaired site would typically be dominated by these more tolerant species, and as a result biodiversity at the site would be quite low.

Threshold values to classify the water quality of watercourses based on Hilsenhoff Family Biotic Index (1987)	
Family Biotic Index	Water Quality Rating
0.00-3.75	Excellent
3.76-4.25	Very Good
4.26-5.00	Good
5.01-5.75	Fair
5.76-6.50	Fairly Poor
6.51-7.25	Poor
7.26-10.00	Very Poor

4.0 SURFACE WATER QUALITY MONITORING RESULTS

The Canadian Council of Ministers of the Environment (CCME) Water Quality Index (WQI) was used to summarize the indicator parameter data collected from NPCA surface water quality monitoring stations between 2018 and 2022. This approach reduces the overall sample size of water quality monitoring stations, but it allows for the partitioning of the water quality dataset to determine if CCME WQI ratings are changing over time. This approach is consistent with Conservation Ontario's recommendation for comparing water quality data in watershed reporting (Conservation Ontario 2011). Using


the five-year blocks of data minimizes seasonal variation and provides sufficient data (n=40 to 50) for reliable statistics in surface water analysis.

The WQI was developed by a sub-committee established under the Canadian Council for Ministers of the Environment (CCME) Water Quality Guidelines Task Group to provide a convenient means of summarizing complex water quality information and communicating it to the public (CCME 2001). The WQI incorporates the number of parameters where water quality objectives have been exceeded, the frequency of exceedances within each parameter, and the amplitude of each exceedance. The index produces a number between 0 and 100 which represents the worst and best water quality, respectively. These numbers are divided into five descriptive categories that range from *poor* to *excellent* (**Table 2**). The CCME WQI has been used extensively by other agencies, including conservation authorities and provincial ministries, as a means of reporting water quality data.

Table 2: CCME Water Quality Index categories (CCME 2001)

CATEGORY	WATER QUALITY INDEX	DESCRIPTION
Excellent	95-100	Water quality is protected with a virtual absence of threat or impairment; conditions very close to natural or pristine levels.
Good	80-94	Water quality is protected with only a minor degree of threat or impairment; conditions rarely depart from natural or desirable levels.
Fair	65-79	Water quality is usually protected but occasionally threatened or impaired; conditions sometimes depart from natural or desirable levels.
Marginal	45-64	Water quality is frequently threatened or impaired; conditions often depart from natural or desirable levels.
Poor	0-44	Water quality is almost always threatened or impaired; conditions usually depart from natural or desirable levels.

The calculation of the WQI is dependent on the water quality parameters and objectives selected for analysis. The indicator parameters and objectives summarized in **Table 1** were used to determine the WQI for NPCA monitoring stations. Benthic invertebrate data is not included in the WQI and is presented separately. It is important to note that the water quality information presented in this report is limited by the size of the dataset which represents 1 to 5 years of data, depending on the station. The reliability of the WQI rating improves over time (> 3 years) as more data is collected and a wider range of water quality conditions are captured in the dataset. In addition, water quality indicator parameters were summarized using boxplots which allow for comparison of the data distributions and basic statistical attributes such as median, percentiles, overall range, and outliers.



Example of a boxplot

4.1 TREND ANALYSIS

The NPCA surface water quality database has reached a sufficient size for trend analysis to be conducted. Trend analyses are very useful for determining if water quality parameter concentrations are increasing, decreasing, or remaining unchanged over time. If the concentration of a water quality parameter is found to be increasing or remaining in an impaired status, then appropriate corrective action can be taken. Trend analysis is also useful for evaluating the performance of stewardship or remediation efforts.

The data on many water quality parameters for the NPCA are not normally distributed and it is not appropriate to use parametric statistical methods to test for trends. Non-parametric statistical methods can deal effectively with non-normally distributed data and are flexible enough to account for seasonal variability. The Seasonal Mann-Kendall Test is often used to determine trends in water quality data (Helsel and Hirsch 1992). The Seasonal Mann-Kendall Test modified from the Mann-Kendall Test (Helsel and Hirsch 1992), compares relative ranks of data values from the same season. This means the water quality parameter concentrations of May would be compared with concentrations of May in other years. Similarly, June concentrations would be compared with June concentrations and so forth. The null hypothesis (H_0) is that the concentration of a water quality parameter is independent of time or, in other words, the datasets show no distinct trend. The alternative hypothesis (H_A) means that a significant increasing or decreasing trend is found over time. The Seasonal Mann-Kendall uses alpha (α) to quantify the probability that a trend exists. For this report, the alpha level for statistical significance was set at $\alpha = 0.05$. This alpha level is commonly used in statistical methods to test for statistical significance. It should be noted that a value of $\alpha = 0.05$ means there is a 5

percent possibility of falsely rejecting the null hypothesis that no trend exists. Probability values of less than 0.05 mean there was statistically significant trend (increasing or decreasing). Trend analysis using the Seasonal Mann-Kendall Test was conducted on chloride, *E. coli*, total phosphorus and total suspended solids concentrations at all stations with 5 or more years of data using software provided by the U.S. Geological Survey (Helsel *et al.*, 2005). Trend analysis for copper, lead, nitrate, and zinc parameters could only be conducted on a small number of stations because many concentrations found were below the laboratory detection limits. These were reported as “non-detect” or a “less than” the laboratory detection limit. Trend analysis with many non-detections or less than values was not favourable for analysis and therefore was excluded from most stations.

4.2 WELLAND RIVER WATERSHED

The Welland River is the largest watershed in the NPCA jurisdiction with a total drainage area of 1,023 km². The watershed covers eleven local municipalities, originating in the Town of Ancaster and spanning the center of the Niagara Peninsula to its physical outlet in the City of Niagara Falls at the Niagara River (**Figure 1**). Over 70% of the watershed is classified as rural. The Welland River is part of the Niagara River Area of Concern (AOC). As shown in **Appendix A**, 30 of the 80 surface water quality monitoring stations are in the Welland River watershed, and 14 of these 30 stations are located on the main Welland River channel.

4.2.1 Welland River: Canadian Water Quality Index

The calculated WQI for the Welland River ranges from *poor* to *fair*. Based on the 2018-2022 data collected, six of fourteen Welland River stations have *poor* water quality, six stations were rated as *marginal*, and two stations were rated as *fair*. WQI results are illustrated in **Appendix A**. Mapping showing the spatial distribution and boxplots of the eight indicator parameters from 2018 to 2022 are found in **Appendix B** to **Appendix I**. Highlights of the water quality monitoring in the Welland River are summarized in **Table 3**:



Figure 1: Map of the subwatersheds monitored for water quality within the Welland River watershed

Table 3: Summary of NPCA water quality data for the Welland River (2018-2022)

STATION	WQI RATING ↔ Stable ↓ Declining ↑ Improving	HILSENHOFF FAMILY BIOTIC INDEX RATING	FACTORS AFFECTING WATER QUALITY (%)= PERCENTAGE OF SAMPLES EXCEEDING GUIDELINES THIS IS ONLY REPORTED WHEN >50% OF SAMPLES EXCEED GUIDELINE	TREND GREEN- DECREASING BLACK- STABLE RED- INCREASING
WR00A Welland River	Marginal ↔	Poor	<ul style="list-style-type: none"> Exceedances of copper, <i>E. coli</i>, lead, total phosphorus (95%), total suspended solids and zinc. Potential stressors include: agricultural and roadway run-off Groundwater discharges sustains continuous baseflow at this site. 	<ul style="list-style-type: none"> Decreasing total phosphorus and total suspended solid concentrations Stable <i>E. coli</i> concentrations Increasing chloride concentrations
WR000 Welland River	Fair ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances of <i>E. coli</i> and total phosphorus (67%) Potential stressors include: agricultural and roadway run-off Groundwater discharge provides intermittent baseflow but the watercourse will dry up in the summer when groundwater levels drop 	<ul style="list-style-type: none"> Decreasing chloride and <i>E.coli</i> concentrations Stable total phosphorus and total suspended solid concentrations
WR001 Welland River	Marginal ↑	Poor	<ul style="list-style-type: none"> Exceedances of chloride, copper, <i>E. coli</i> (58%), total phosphorus (77%), total suspended solids and zinc Potential stressors include: agricultural, airport and roadway run-off 	<ul style="list-style-type: none"> Decreasing zinc concentrations Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
WR002 Welland River	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances of chloride (93%), copper (55%), <i>E. coli</i>, lead, total phosphorus, total suspended solids and zinc (90%) Potential stressors include: agricultural, airport and roadway run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus total suspended solid, and zinc concentrations
WR020 Welland River	Marginal ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedances in chloride (73%), copper, <i>E. coli</i>, total phosphorus (95%), and total suspended solids Potential stressors include agricultural and roadway run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus total suspended solid, and zinc concentrations
WR003 Welland River	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances of chloride (56%), copper, <i>E. coli</i>, nitrate, total phosphorus (97%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Decreasing total suspended solid concentrations Stable <i>E. coli</i> and total phosphorus concentrations Increasing chloride concentrations

WR004 Welland River	Fair ↑	Very Poor	<ul style="list-style-type: none"> Exceedances of <i>E. coli</i>, total phosphorus (81%) and total suspended solids. Potential stressors include: agricultural and roadway run-off Lake Niapenco is improving the water quality the Welland River at this site 	<ul style="list-style-type: none"> Decreasing <i>E.coli</i>, total phosphorus and total suspended solid concentrations Stable chloride concentrations
WR005 Welland River	Poor ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances of chloride, copper, <i>E. coli</i> (57%), nitrate, total phosphorus (100%), suspended solids (74%) and zinc. Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E.coli</i>, total phosphorus and suspended solid concentrations
WR006 Welland River	Marginal ↑	Poor	<ul style="list-style-type: none"> Exceedances of copper, <i>E. coli</i>, nitrate, total phosphorus (100%), suspended solids and zinc Potential stressors include: agricultural and roadway run-off Algae and duckweed observed during summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
WR007 Welland River	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances of copper, <i>E. coli</i>, nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural, roadway run-off Algae and duckweed observed during summer months Site is invaded by non-native Zebra Mussels 	<ul style="list-style-type: none"> Stable chloride and total phosphorus concentrations Increasing <i>E. coli</i> and total suspended solid concentrations
WR009B Welland River	Marginal ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedances of copper, <i>E. coli</i> (51%), total phosphorus (95%), total suspended solid and zinc Potential stressors include: Sewage treatment plant effluent and agricultural and urban run-off Site strongly influenced by Niagara River backwater which has the potential to improve water quality 	<ul style="list-style-type: none"> Decreasing chloride concentrations Stable <i>E. coli</i>, total phosphorus and total suspended solid concentrations
WR010 Welland River	Marginal ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedances of copper, <i>E. coli</i>, total phosphorus (93%), total suspended solids and zinc. Potential stressors include: Sewage treatment plant effluent and agricultural and urban run-off Site strongly influenced by Niagara River backwater which has the potential to improve water quality 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total phosphorus concentrations
WR011 Welland River	Marginal ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedances of copper, <i>E. coli</i>, total phosphorus (79%) and total suspended solids Potential stressors include: Sewage treatment plant effluent and agricultural and urban run-off Site strongly influenced by Niagara River backwater which has the potential to improve water quality 	<ul style="list-style-type: none"> Decreasing <i>E. coli</i> concentrations Stable total phosphorus and total suspended solid concentrations Increasing chloride concentrations

WR012 Welland River	Fair ↔	Insufficient Data	<ul style="list-style-type: none"> • Exceedances of <i>E. coli</i>, total phosphorus and total suspended solids • Potential stressors include: Urban run-off • Site strongly influenced by Niagara River backwater which has the potential to improve water quality 	<ul style="list-style-type: none"> • Stable chloride, <i>E. coli</i>, total phosphorus and total phosphorus concentrations
------------------------	-----------	----------------------	--	---



Duckweed Bloom on the Welland River in West Lincoln

4.2.2 WELLAND RIVER: HILSENHOFF BIOTIC INDEX RESULTS

Hilsenhoff Biotic Index (HBI) results indicate that water quality at most stations in the Welland River ranged from *Very Poor* to *Poor* (**Table 3**). Two sites managed to obtain a *Fairly Poor* rating (WR000 and WR005). Results from Hilsenhoff Biotic Index assessments completed between 2018 and 2022 are illustrated in **Appendix J**.

Low HBI scores observed in the Welland River mainly are due to road salts and metals in stormwater, sediment loading, lack of in-stream habitat, and nutrient enrichment. A biological assessment was not completed for WR009B, WR010, WR011 and WR012 due to high water depth and channel morphology. These stations are located at the siphon where the Welland River flows beneath the Welland Canal and would require boat access for sample collection.

4.2.3 WELLAND RIVER: KEYS FINDINGS

- Based on the 2018-2022 data, elevated concentrations of total phosphorus are a widespread cause of water quality impairment in the Welland River. Greater than 95% of samples collected in the main Welland River exceeded the PWQO with some concentrations greater than 20 times the PWQO. High phosphorus in the Welland River has stimulated the overgrowth of algae and duckweed throughout the watershed. When these plants transpire, and decompose they deplete dissolved oxygen in the water and this in turn stresses aquatic organisms such as fish and benthic invertebrates. Manure from livestock operations, sewage discharges, soil erosion, fertilizers, and pesticides are sources of total phosphorus in the Welland River.
- Generally, the overall water quality of the Welland River downstream of the City of Welland is less stressed than the water upstream of the City of Welland. This is caused by the redirection of the Niagara River water down the Welland River in Chippawa for Ontario Power Generation (OPG). This results in a dilution effect that reduces the concentrations of water quality parameters. This effect is observed to the east side of the City of Welland. However, upstream of the City of Welland, the river flow pattern caused by OPG operations and canal siphons are likely restricting the natural flushing of sediment, nutrients and other contaminants from the central Welland River watershed and exacerbating water quality conditions in this watershed.
- Water quality stations in the vicinity of Hamilton Airport (HIA) continue to have water quality designated as *poor* due to elevated concentrations of chloride and zinc. Chloride concentrations are stable at WR001 but increasing at WR002 despite the recent removal of the road salt storage pad. Zinc concentrations found at these stations consistently exceed the PWQO and are the highest observed in the NPCA water quality network. The current information that the HIA has suggests that zinc is coming off the brake system of the airplanes. It should be noted that zinc concentrations have been decreasing at both stations. The NPCA also has not observed any propylene glycol discharge in WR001 or WR002 this year. In 2011, the HIA expanded its facilities and upgraded its water quality safeguards to WR001 and WR002. Continued monitoring by the NPCA will track water quality changes at these tributaries. The NPCA does not monitor the water quality of the Hamilton Airport tributary identified as the potential source of Per- and polyfluoroalkyl substances (PFAS) that has been found in

turtle/fish tissue sampled at Binbrook Conservation Area. PFAS are a man-made compound belonging to a large family of compounds known as perfluorinated chemicals. These compounds do not readily breakdown and have the potential to bioaccumulate in animal tissue. MECP continues to provide fish consumption guidelines based on fish samples they have collected for this area and information is found on (<https://www.ontario.ca/page/guide-eating-ontario-fish>). The NPCA continues to notify Binbrook Conservation Area Park users about the new fish consumption guidelines and information regarding PFAS has been posted on the NPCA website: <https://npca.ca/parks-recreation/conservation-areas/binbrook>. Since 2015, Transport Canada and Procurement Canada have retained Arcadis Canada Inc. to conduct a risk assessment to investigate presence and distribution of PFAS in the Welland River downstream of the HIA. Through this assessment process Arcadis has released project updates to property owners and other groups with an interest in the risk assessment area. The final report is still pending. The NPCA Watershed Monitoring and Reporting division has added PFAS sampling in 2012 as part of special project monitoring program at Binbrook Reservoir and this information can be found in Section 6.4.

4.3 WELLAND RIVER TRIBUTARIES

Fifteen tributaries of the Welland River are monitored through the NPCA Water Quality Monitoring Program. These tributaries include Buckhorn Creek, Elsie Creek, Mill Creek, Oswego Creek, Beaver Creek, Big Forks Creek, Coyle Creek, Drapers Creek, Feeder Canal, Mill Race Creek, Grassy Brook, Tee Creek, Thompson Creek, Power Canal, and Lyons Creek (**Figure 1**). Tributaries were selected based on drainage area, landuse, restoration projects, and watershed plans.

4.3.1 WELLAND RIVER TRIBUTARIES: WATER QUALITY INDEX

Based on the results of the WQI fifteen of sixteen Welland River tributary stations have water quality that is rated as *poor* (**Table 4**). The Power Canal (PR001) was found to have water quality rated as *marginal*. WQI results are illustrated in **Appendix A**. Mapping showing the spatial distribution and boxplots of the eight indicator parameters from 2018 to 2022 are found in **Appendix B** to **Appendix I**. Highlights of the water quality monitoring in the Welland River are summarized in **Table 4**:

Table 4: Summary of NPCA water quality data for Welland River tributaries (2018-2022)

STATION WATERSHED	WQI RATING ↔ Stable ↓ Declining ↑ Improving	HILSENHOFF FAMILY BIOTIC INDEX RATING	FACTORS AFFECTING WATER QUALITY (%)= PERCENTAGE OF SAMPLES EXCEEDING GUIDELINES THIS IS ONLY REPORTED WHEN >50% OF SAMPLES EXCEED GUIDELINE	TREND GREEN- DECREASING BLACK- NO TREND RED- INCREASING
BF001 Big Forks Creek	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (44%), nitrate, total phosphorus (100%), total suspended solids and zincs. Potential stressors include: agricultural and roadway run-off Significant algae and overabundance of duckweed observed during summer months Prone to zero baseflow conditions in the summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, and total suspended solid concentrations Increasing total phosphorus concentrations
BU001 Buckhorn Creek	Poor ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride (51%), copper, <i>E. coli</i> (64%), lead, nitrate, total phosphorus (95%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Prone to zero baseflow conditions in the summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
BV001 Beaver Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i>, lead, nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Significant algae observed during summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
CO001 Coyle Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i>, nitrate, total phosphorus (97%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Site invaded by non-native Zebra Mussels 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable <i>E. coli</i> and total phosphorus concentrations Increasing chloride concentrations
DR001 Drapers Creek	Poor ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (92%), total phosphorus (100%), and zinc Potential stressors include: urban run-off Algae observed during summer months 	<ul style="list-style-type: none"> Stable total phosphorus and total suspended solid concentrations Increasing <i>E. coli</i> and chloride concentrations

EL001 Elsie Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i>, lead, nitrate, total phosphorus (97%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Algae observed during summer months Prone to zero baseflow conditions in the summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus, and total suspended solid concentrations
GR001 Grassy Brook	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (53%), lead, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Algae observed during summer months Prone to zero baseflow conditions in the summer months 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable chloride, <i>E. coli</i> and total phosphorus concentrations
TE001 Tee Creek	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (54%), nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Nutrient enrichment from upstream agricultural areas Prone to zero baseflow conditions in the summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
LY003 Lyons Creek	Marginal ↑	Poor	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i>, lead, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Site strongly influenced by Niagara River backwater which has the potential to improve water quality 	<ul style="list-style-type: none"> Decreasing chloride and total phosphorus concentrations Stable <i>E. coli</i> and total suspended solid concentrations
MI001 Mill Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i>, lead, nitrate, total phosphorus (97%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Algae and overabundance of duckweed observed during summer months 	<ul style="list-style-type: none"> Decreasing total phosphorus and total suspended solid concentrations Stable <i>E. coli</i> concentrations Increasing chloride concentrations
OS001 Oswego Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i> (58%), nitrate, total phosphorus (100%), total suspended solids (62%) and zinc Potential stressors include: agricultural and roadway run-off Algae and overabundance of duckweed observed during summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i> and total suspended solids concentrations Increasing total phosphorus

OS002 Oswego Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper (51%), <i>E. coli</i> (66%), lead, nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Algae and overabundance of duckweed observed during summer months 	<ul style="list-style-type: none"> Stable <i>E. coli</i>, total phosphorus and total suspended solid concentrations Increasing chloride concentrations
TC001 Thompson Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (65%), lead, nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Stable <i>E. coli</i>, total phosphorus and total suspended solid concentrations Increasing chloride concentrations
PR001 Power Canal	Marginal ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedances in total phosphorus, total suspended solids and zinc. Potential stressors include: urban run-off and Niagara Falls waste water treatment plant Water source at this site is Niagara River water which potentially improves water quality 	<ul style="list-style-type: none"> Decreasing <i>E. coli</i> concentrations Stable chloride, total phosphorus and total suspended solid concentrations
MR001 Mill Race Creek	Poor ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i> (66%), lead, nitrate, total phosphorus (100%) and total suspended solids and zinc. Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
FC001 Feeder Canal	Poor ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i>, lead, total phosphorus (88%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Decreasing <i>E. coli</i> and chloride concentrations Stable total phosphorus and total suspended solid concentrations

4.3.2 WELLAND RIVER TRIBUTARIES: HILSENHOFF BIOTIC INDEX RESULTS

HBI results indicate that water quality is ranged from *very poor* to *fairly poor* at all Welland River tributary stations currently monitored (**Table 4**). Results from biological assessments completed between 2018 and 2022 are illustrated in **Appendix J**. Generally, the HBI results match with water chemistry ratings. Sediment loading, lack of in-stream habitat, and nutrient enrichment are the primary causes of impairment at all stations.

4.3.3 WELLAND RIVER TRIBUTARIES: KEY FINDINGS

- Based on the 2018-2022 data, elevated concentrations of total phosphorus are a widespread cause of water quality impairment in the Welland River tributaries. Approximately 95% of samples collected from the Welland River tributaries exceeded the PWQO with some concentrations greater than 30 times the PWQO. Concentrations of total phosphorus are very high in Beaver Creek, Big Forks Creek, Oswego Creek, and Tee Creek. It is recommended that these subwatersheds be prioritized by Best Management Practice programs such as those provided by the NPCA to reduce phosphorus loads. Sources of phosphorus include manure from livestock operations, sewage discharges, soil erosion, fertilizers, and pesticides.
- *E. coli* concentrations frequently exceed the PWQO in Buckhorn Creek, Big Forks Creek, Beaver Creek, Coyle Creek, Drapers Creek, Elsie Creek, Mill Creek, and Oswego Creek.
- Decreasing total phosphorus concentrations in Mill Creek are now being observed with the NPCA's long-term data. Mill Creek watersheds have been targeted by the NPCA's stewardship program over the last twenty years to reduced non-point pollution such as nutrients and sediment. This data provides some evidence that this program is reducing nutrient runoff.



NPCA water quality sensors

4.4 TWENTY MILE CREEK WATERSHED

The Twenty Mile Creek watershed is the second largest watershed in the NPCA jurisdiction with a total drainage area of 302 km². Nine of 80 NPCA surface water quality monitoring stations are located within the Twenty Mile Creek watershed. There are six stations on the main channel. There are also monitoring stations for each of the subwatersheds which include Spring Creek, North Creek and Gavora Ditch (**Figure 4**).




Figure 4: Map of the subwatersheds monitored for water quality within the Twenty Mile Creek watershed

4.4.1 TWENTY MILE CREEK WATERSHED: CANADIAN WATER QUALITY INDEX

Based on the results of the WQI five of nine Twenty Mile Creek watershed stations have water quality that is rated as *poor*. Four stations (GV001, SP001, TN002 and TN003) were rated as *marginal*. WQI results are illustrated in **Appendix A**. Mapping showing the spatial distribution and boxplots of the eight indicator parameters from 2018 to 2022 are found in **Appendix B** to **Appendix I**. Highlights of the water quality monitoring in the Twenty Mile Creek are summarized in **Table 5**:

Table 5: Summary of NPCA water quality data for the Twenty Mile Creek watershed (2018-2022)

STATION WATERSHED	WQI RATING ↔ Stable ↓ Declining ↑ Improving	HILSENHOFF FAMILY BIOTIC INDEX RATING	FACTORS AFFECTING WATER QUALITY (%) = PERCENTAGE OF SAMPLES EXCEEDING GUIDELINES THIS IS ONLY REPORTED WHEN >50% OF SAMPLES EXCEED GUIDELINE	TREND GREEN- DECREASING BLACK- NO TREND RED- INCREASING
TN001 Twenty Mile Creek	Poor ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (54%), lead, total phosphorus (92%), total suspended solids and zinc Potential stressors include: agricultural and urban run-off Site invaded by the non-native Chinese Mystery Snails Excessive algae observed during the summer months 	<ul style="list-style-type: none"> Decreasing total phosphorus concentrations Stable chloride, <i>E. coli</i> and total suspended solid concentrations
TN002 Twenty Mile Creek	Marginal ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i>, total phosphorus (90%), and total suspended solids. Potential stressors include: agricultural and urban run-off Prone to zero baseflow conditions in the summer months 	<ul style="list-style-type: none"> Decreasing total phosphorus concentrations Stable chloride, <i>E. coli</i> and total suspended solid concentrations
TN003 Twenty Mile Creek	Marginal ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i>, nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and urban run-off Excessive algae observed during the summer months 	<ul style="list-style-type: none"> Decreasing total suspended solid concentrations Stable chloride, <i>E. coli</i>, and total phosphorus concentrations
TN003A Twenty Mile Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (50%), nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and urban run-off Excessive algae observed during the summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus, and total suspended solid concentrations
TN004 Twenty Mile Creek	Poor ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (68%), nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Algae observed during the summer months 	<ul style="list-style-type: none"> Stable <i>E. coli</i>, total phosphorus and total suspended solid concentrations Increasing chloride concentrations

TN006 Twenty Mile Creek	Poor ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i>, nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Excessive algae observed during the summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus, and total suspended solid concentrations
NC001 North Creek	Poor ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (74%), nitrate, total phosphorus (97%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off Prone to zero baseflow conditions in the summer months Excessive algae observed during the summer months 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable <i>E. coli</i> and total phosphorus concentrations Increasing chloride concentrations
SP001 Spring Creek	Poor ↓	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (65%), lead, total phosphorus (100%) and zinc Potential stressors include: agricultural and roadway run-off Prone to zero baseflow conditions in the summer months Excessive algae observed during the summer months 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable <i>E. coli</i> and total phosphorus concentrations Increasing chloride concentrations
GV001 Gavora Ditch	Marginal ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i> (54%), and total phosphorus (100%) Potential stressors include: agricultural and roadway run-off Prone to zero baseflow conditions in the summer months Algae observed during summer months 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable chloride, <i>E. coli</i>, and total phosphorus concentrations

4.4.2 TWENTY MILE CREEK WATERSHED: HILSENHOFF BIOTIC INDEX RESULTS

HBI results indicate that water quality is ranged from *poor* to *fairly poor* at most Twenty Mile Creek monitoring stations (**Table 5**). Results from biological assessments completed between 2018 and 2022 are illustrated in **Appendix J**. Reduced baseflow, high sediment loading due to erosion, lack of in-stream habitat, and nutrient enrichment are primary causes of impairment at these stations.

4.4.3 TWENTY MILE CREEK WATERSHED: KEY FINDINGS

- Based on the 2018-2022 data, elevated concentrations of total phosphorus are a widespread cause of water quality impairment in the Twenty Mile watershed. Approximately 95% of samples collected from the Twenty Mile watershed exceeded the PWQO with some concentrations greater than 30 times the PWQO.
- *E. coli* and total suspended solid concentrations frequently exceed the provincial objective in Twenty Mile Creek watershed. It is recommended that this subwatershed be prioritized by Best Management Practice programs such as those provided by the NPCA to reduce sources of *E. coli* in this watershed.




NPCA staff collecting water sampling from Oswego Creek.

4.5 LAKE ONTARIO TRIBUTARIES

Nineteen tributaries discharging into Lake Ontario are monitored through the NPCA Water Quality Monitoring Program. These tributaries include: Forty Mile Creek, Thirty Mile Creek, Eighteen Mile Creek, Sixteen Mile Creek, Fifteen Mile Creek, Twelve Mile Creek, Francis Creek, Richardson Creek, Walker's Creek, Eight Mile Creek, Six Mile Creek, Four Mile Creek, Two Mile Creek, One Mile Creek, Bartlett Creek, Purdhommes Drain, Welland Canal, Shriners Creek, and Beaver Dam Creek (**Figure 6**). Twenty Mile Creek is also a tributary of Lake Ontario but is presented separately due to the relatively large size of the watershed.

4.5.1 LAKE ONTARIO TRIBUTARIES: CANADIAN WATER QUALITY INDEX

Based on the results of the WQI, eighteen of twenty-seven Lake Ontario tributary stations have water quality that is rated as *poor*. Seven stations were rated as *marginal*, and two stations were rated as *fair*. WQI results are illustrated in **Appendix A**. Mapping showing the spatial distribution and boxplots of the eight indicator parameters from 2018 to 2022 are found in **Appendix B** to **Appendix I**. Highlights of the water quality monitoring in the Lake Ontario tributaries are summarized in **Table 6**.



Twenty Mile Creek station in Jordan




Figure 6: Map of the subwatersheds draining to Lake Ontario that are monitored for water quality

Table 6: Summary of NPCA water quality data for Lake Ontario tributaries (2018-2022)

STATION WATERSHED	WQI RATING ↔ Stable ↓ Declining ↑ Improving	HILSENHOFF FAMILY BIOTIC INDEX RATING	FACTORS AFFECTING WATER QUALITY (%)= PERCENTAGE OF SAMPLES EXCEEDING GUIDELINES THIS IS ONLY REPORTED WHEN >50% OF SAMPLES EXCEED GUIDELINE	TREND GREEN- DECREASING BLACK- NO TREND RED- INCREASING
FM001 Forty Mile Creek	Poor ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride (60%), copper, <i>E. coli</i> (85%), total phosphorus (98%), total suspended solids and zinc Potential stressors include: road salt storage compound, quarry dewatering, urban and agricultural run-off. Algae observed during summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
ET001 Eighteen Mile Creek	Poor ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (64%), nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: greenhouse waste water, rural and agricultural run-off. Very frequent copper exceedances warrant further investigation 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
FF001 Fifteen Mile Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (54%), lead, total phosphorus (100%), total suspended solids and zinc Excessive algae observed during summer months Potential stressors include: rural and agricultural run-off 	<ul style="list-style-type: none"> Decreasing total suspended solid concentrations Stable chloride, <i>E. coli</i>, and total phosphorus concentrations.
SX001 Sixteen Mile Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (72%), lead, nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
EI001 Eight Mile Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (84%), nitrate, total phosphorus (100%), and total suspended solids and zinc. Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Stable <i>E. coli</i> and total suspended solid concentrations Increasing chloride and total phosphorus concentrations

FA001 Francis Creek	Marginal ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride (79%), copper, <i>E. coli</i> (71%), total phosphorus (68%), total suspended solids and zinc. Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Stable <i>E. coli</i>, total phosphorus and total suspended solid concentrations Increasing chloride concentrations
RC001 Richardson Creek	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride (77%), copper (65%), <i>E. coli</i> (56%), nitrate (100%), and total phosphorus (100%), total suspended solids and zinc. Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Stable <i>E. coli</i> and total suspended solid concentrations Increasing chloride and total phosphorus concentrations
SI001 Six Mile Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride (59%), copper, <i>E. coli</i> (72%), nitrate, total phosphorus (79%), total suspended solids and zinc Potential stressors include: agricultural and roadway run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
FU004 Four Mile Creek	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (69%), nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and rural run-off 	<ul style="list-style-type: none"> Decreasing <i>E. coli</i> and total suspended solid concentrations Stable chloride concentrations Increasing total phosphorus concentrations
TM001 Two Mile Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (87%), nitrate, total phosphorus (100%), and total suspended solids Excessive <i>E. coli</i> concentrations warrant further investigations Potential stressors include: rural and urban run-off 	<ul style="list-style-type: none"> Decreasing chloride and total suspended solid concentrations Stable <i>E. coli</i> and total phosphorus concentrations
OM001 One Mile Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride (56%), copper, <i>E. coli</i> (94%), nitrate, total phosphorus (100%), total suspended solids and zinc Potential stressors include: urban run-off Prone to zero baseflow conditions in the summer months 	<ul style="list-style-type: none"> Stable <i>E. coli</i> and total suspended solid concentrations Increasing chloride and total phosphorus concentrations

TW001 Twelve Mile Creek	Marginal ↑	Fairly Poor	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i> (56%), total phosphorus (61%) and total suspended solids. Potential stressors include: rural and urban run-off Groundwater discharges sustains continuous baseflow at this site. 	<ul style="list-style-type: none"> Decreasing <i>E. coli</i> and total suspended solid concentrations. Stable total phosphorus concentrations Increasing chloride concentrations
TW002 Twelve Mile Creek	Fair ↑	Fairly Poor	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i>, total phosphorus, and total suspended solids. Potential stressors include: agricultural and rural run-off Groundwater discharges sustains continuous baseflow at this site. 	<ul style="list-style-type: none"> Decreasing total phosphorus and total suspended solids Stable chloride and <i>E. coli</i> concentrations
TW003 Twelve Mile Creek	Marginal ↑	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (62%), total phosphorus (89%), total suspended solids and zinc Potential stressors include: decommissioned landfill and rural run-off Groundwater discharges sustains continuous baseflow at this site. 	<ul style="list-style-type: none"> Stable <i>E. coli</i>, total phosphorus and total suspended solid concentrations Increasing chloride concentrations
TW004 Twelve Mile Creek	Marginal ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i>, nitrate (97%), total phosphorus, and total suspended solids Potential stressors include: golf course and rural run-off Groundwater discharges sustains continuous baseflow at this site. 	<ul style="list-style-type: none"> Decreasing total phosphorus concentrations Stable <i>E. coli</i> and total suspended solid concentrations Increasing chloride and nitrate concentrations
TW005 Twelve Mile Creek	Marginal ↔	Fair	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (59%), total phosphorus (73%), and total suspended solids Potential stressors include: rural and urban run-off Groundwater discharges sustains continuous baseflow at this site. 	<ul style="list-style-type: none"> Decreasing <i>E. coli</i> and total suspended solids concentrations Stable total phosphorus concentrations Increasing chloride concentrations
TW006 Twelve Mile Creek	Fair ↔	Fair	<ul style="list-style-type: none"> Exceedances in <i>E. coli</i> (54%), total phosphorus (64%) and total suspended solids Potential stressors include: rural run-off Groundwater discharges sustains continuous baseflow at this site. 	<ul style="list-style-type: none"> Decreasing chloride and <i>E. coli</i> concentrations Stable total phosphorus and total suspended solid concentrations

TW007 Twelve Mile Creek	Marginal ↔	Fairly Poor	<ul style="list-style-type: none"> Exceedances in chloride, <i>E. coli</i> (67%), nitrate, total phosphorus (81%), and total suspended solids Potential stressors include: agricultural and rural run-off Groundwater discharges sustains continuous baseflow at this site. 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i> and total suspended solid concentrations Increasing total phosphorus concentrations
TW008 Twelve Mile Creek	Marginal ↑	Poor	<ul style="list-style-type: none"> Exceedances in chloride (79%), <i>E. coli</i> (59%), nitrate, and total phosphorus (100%) Potential stressors include: agricultural and rural run-off Prone to zero baseflow conditions in the summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i> and total suspended solids concentrations Increasing total phosphorus concentrations
TW009 Twelve Mile Creek	Marginal ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedances in chloride, <i>E. coli</i> (50%), total phosphorus and total suspended solids Potential stressors include: urban run-off and industrial waste water Water source at this site is predominately from the Welland Canal water which potentially improves water quality 	<ul style="list-style-type: none"> Stable chloride, total phosphorus and total suspended solid concentrations Increasing <i>E. coli</i> concentrations
TH001 Thirty Mile Creek	Marginal ↑	Poor	<ul style="list-style-type: none"> Exceedances in chloride (56%), copper, <i>E. coli</i> (77%), nitrate, total phosphorus (92%) and total suspended solids Potential stressors include: agricultural and rural run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
WC001 Walkers Creek	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride (68%), copper, <i>E. coli</i> (84%), lead, nitrate, total phosphorus (95%), total suspended solids and zinc Potential stressors include: urban run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
SH002 Shriners Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride (69%), copper, <i>E. coli</i>, total phosphorus (100%) and total suspended solids Potential stressors include: urban run-off Algae and duckweed observed during summer months 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable <i>E. coli</i> concentrations Increasing chloride and total phosphorus


BE004 Beaver Dam Creek	Marginal ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, <i>E. coli</i>, total phosphorus (84%), total suspended solids and zinc. Potential stressors include: industrial and urban run-off Algae and duckweed observed during summer months 	<ul style="list-style-type: none"> Decreasing <i>E. coli</i> and total phosphorus concentrations Increasing chloride and total suspended solid concentrations
WE001 Welland Canal	Marginal ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedance in chloride, <i>E. coli</i>, total phosphorus, total suspended solids and zinc. Water source at this site is predominately from the Lake Erie 	<ul style="list-style-type: none"> Stable <i>E. coli</i>, total phosphorus and total suspended solid concentrations Increasing chloride concentrations
PD001 Prudhommes Drain	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride (80%), copper, <i>E. coli</i> (83%), lead, nitrate, total phosphorus (93%), total suspended solids and zinc Potential stressors include: urban run-off Algae and duckweed observed during summer months 	<ul style="list-style-type: none"> Decreasing total phosphorus and total suspended solids Stable <i>E. coli</i> concentrations Increasing chloride concentrations
BT001 Bartlett Creek	Poor ↓	Poor	<ul style="list-style-type: none"> Exceedances in chloride (61%), copper, <i>E. coli</i>, nitrate, total phosphorus (96%), total suspended solids and zinc Potential stressors include: highway and agricultural run-off 	<ul style="list-style-type: none"> Stable <i>E. coli</i>, total phosphorus and total suspended solid concentrations Increasing chloride concentrations

4.5.2 LAKE ONTARIO TRIBUTARIES: HILSENHOFF BIOTIC INDEX RESULTS

HBI results indicate that water quality is ranged from *fair* to *very poor* at Lake Ontario tributary stations (**Table 6**). Results from biological assessments completed between 2018 and 2022 are illustrated in **Appendix J**. Sediment loading, nutrient enrichment, and the lack of in-stream habitat are the primary causes of impairment at these stations. Upper Twelve Mile Creek station TW006 located on the Effingham tributary are rated as *fair*. The Effingham tributary of upper Twelve Mile Creek is the only watercourse in the NPCA watershed that consistently achieves this rating. This site can support several sensitive taxa such as mayflies and stoneflies due to cooler water temperatures, excellent riparian buffer and in-stream habitat, and suitable water quality.

4.5.3 LAKE ONTARIO TRIBUTARIES: KEY FINDINGS

- The Upper Twelve Mile Creek watershed represents some of the best water quality in the Niagara Peninsula. This portion of Twelve Mile Creek supports brook trout and a rich macroinvertebrate community that is unique in Niagara. The main stresses to the aquatic community include exceedances of total phosphorus and *E. coli*. Nitrate contamination has been identified as a stressor at TW004 and it may be sourced to a local business. Efforts to minimize these stressors through BMP initiatives will allow this watershed to remain in its current state.
- Based on the 2018-2022 data, all the Lake Ontario tributaries have total phosphorus exceedances. The most impacted of these tributaries include Fifteen Mile Creek, Sixteen Mile Creek and Eighteen Mile Creek which had median concentrations nearly 7 times the PWQO. The upper portions of these watersheds need to be prioritized for BMPs to reduce phosphorus loads. Total phosphorus concentrations were found to be lower in the NOTL watersheds.
- Two Mile Creek (TM001) has the highest concentrations of *E. coli* in the NPCA watershed and continues to suggest that there may be sewage entering the Two Mile Creek. The NPCA sampled the stormwater outfalls of Two Mile Creek Conservation Area and found two sources of the bacteria and are currently working with the Town of NOTL and MECP to solve this issue. Elevated *E.coli* concentrations for Walkers Creek (St. Catharines) and Purdhommes Drain (Lincoln) were also observed and additional DNA analysis will be initiated to source of the *E.coli* (animals vs human).
- Copper and zinc concentrations in Eighteen Mile Creek consistently exceed PWQOs. Within the NPCA water quality monitoring network regular metal exceedances are uncommon but based on the landuse in this watershed there may be pesticides entering the watercourse. The NPCA will investigate these exceedances further.
- Richardson Creek consistently exceeds CCME for nitrate and a significant upstream source exists that requires further investigation.



Water Penny from Forty Mile Creek in Grimsby

4.6 NIAGARA RIVER TRIBUTARIES

Four tributaries discharging to the Niagara River are monitored through the NPCA Water Quality Monitoring Program. These tributaries include: Bayer Creek, Black Creek (including Beaver Creek), Frenchman's Creek, and Usshers Creek (**Figure 8**).

4.6.1 NIAGARA RIVER TRIBUTARIES: CANADIAN WATER QUALITY INDEX

Based on the results of the WQI, Usshers Creek (US001), Bayer Creek (BA001), and Frenchman Creek (FR003) stations were rated as *poor* water quality, Black Creek (BL003) station was rated as *marginal* and Beaver Creek (BR001) was rated as *fair*. WQI results are illustrated in **Appendix A**. Mapping showing the spatial distribution and boxplots of the eight indicator parameters from 2018 to 2022 are found in **Appendix B** to **Appendix I**. Highlights of the water quality monitoring in the Niagara River Tributaries are summarized in **Table 7**.



Figure 8: Map of the subwatersheds monitored for water quality in the Niagara River watershed outside of the Welland River

Table 7: Summary of NPCA water quality data for Niagara River tributaries (2018-2022)

STATION WATERSHED	WQI RATING ↔ Stable ↓ Declining ↑ Improving	HILSENHOFF FAMILY BIOTIC INDEX RATING	FACTORS AFFECTING WATER QUALITY (%)= PERCENTAGE OF SAMPLES EXCEEDING GUIDELINES THIS IS ONLY REPORTED WHEN >50% OF SAMPLES EXCEED GUIDELINE	TREND GREEN- DECREASING BLACK- STABLE RED- INCREASING
BA001 Bayer Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (62%), lead, nitrate, total phosphorus (100%), total suspended solids and zinc. Potential stressors include: agricultural and rural run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
BL003 Black Creek	Marginal ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i> (53%), total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and rural run-off 	<ul style="list-style-type: none"> Stable chloride and total phosphorus concentrations Increasing <i>E. coli</i> and total suspended solid concentrations
BR001 Beaver Creek	Fair ↔	Poor	<ul style="list-style-type: none"> Exceedances in <i>E. coli</i>, total phosphorus (97%) and total suspended solids Potential stressors include: agricultural and rural run-off 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable chloride, <i>E. coli</i>, and total phosphorus concentrations
FR003 Frenchman Creek	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (59%), total phosphorus (78%), total suspended solids and zinc Potential stressors include: agricultural and rural run-off Algae observed during summer months 	<ul style="list-style-type: none"> Decreasing chloride and total phosphorus concentrations Stable <i>E. coli</i> and total suspended solid concentrations
US001 Usshers Creek	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i>, total phosphorus (97%), total suspended solids and zinc Potential stressors include: agricultural and rural run-off Prone to zero baseflow conditions in the summer months Algae and duckweed observed during summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations

4.6.2 NIAGARA RIVER TRIBUTARIES: HILSENHOFF BIOTIC INDEX RESULTS

HBI ratings ranged from *very poor* to *poor* at three Niagara River tributary stations (**Table 7**). Results from biological assessments completed between 2018 and 2022 are illustrated in **Appendix J**. Sediment loading, reduced baseflow, lack of in-stream habitat, and nutrient enrichment are primary causes of impairment at these stations. Samples have not been collected from station BL003 due to high water depth, channel morphology, and access restrictions.

4.6.3 NIAGARA RIVER TRIBUTARIES: KEY FINDINGS

- The water quality ratings in Black Creek watershed (BL003 and BR001) were found to be higher than other NPCA watersheds. The degree of landuse impacts from urban and rural pressures are significantly less in this watershed.
- Based on the 2018-2022 data, all the Niagara River tributaries had total phosphorus exceedances. The most impacted of these tributaries was Usshers Creek which had median concentrations 6 times the PWQO. Total phosphorus concentrations were found to be much lower in Frenchman Creek with median concentrations only 2 times the PWQO. Nonetheless these watersheds would benefit by BMP works to reduce phosphorus loads.

4.7 LAKE ERIE TRIBUTARIES

Eight tributaries discharging to Lake Erie are monitored through the NPCA Water Quality Monitoring Program. These tributaries include: Beaver Dam Creek, Casey Drain, Eagle Marsh Drain, Krafts Drain, Low Banks Drain, Point Abino Drain, Six Mile Creek, and Wignell Drain (**Figure 10**). In addition, the Welland Canal monitoring point in Port Colborne is also included with the Lake Erie tributaries. Water at this site enters the Welland Canal and outlets in Lake Ontario.

4.7.1 LAKE ERIE TRIBUTARIES: CANADIAN WATER QUALITY INDEX

Based on the results of the WQI five of nine Lake Erie tributary stations are rated as having *poor* water quality two stations are rated as *marginal*, one station (PA001) rated as fair and one station (WE000) rated as *good* (**Table 8**). WQI results are illustrated in **Appendix A**. Mapping showing the spatial distribution and boxplots of the eight indicator parameters from 2018 to 2022 are found in **Appendix B** to **Appendix I**. Highlights of the water quality monitoring in the Lake Erie Tributaries are summarized in **Table 8**.




Figure 10: Map of the subwatersheds monitored for water quality along the north shore of Lake Erie

Table 8: Summary of NPCA water quality data for Lake Erie tributaries (2018-2022).

STATION WATERSHED	WQI RATING ↔ Stable ↓ Declining ↑ Improving	HILSENHOFF FAMILY BIOTIC INDEX RATING	FACTORS AFFECTING WATER QUALITY (%)= PERCENTAGE OF SAMPLES EXCEEDING GUIDELINES THIS IS ONLY REPORTED WHEN >50% OF SAMPLES EXCEED GUIDELINE	TREND GREEN- DECREASING BLACK- NO TREND RED- INCREASING
BD001 Beaver Dam Drain	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (56%), nickel, total phosphorus (100%), total suspended solids and zinc Potential stressors include: historic industrial pollution, agricultural and rural run-off 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations
CD001 Casey Drain	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (68%), nitrate, total phosphorus (97%), total suspended solids and zinc Potential stressors include: agricultural and rural run-off 	<ul style="list-style-type: none"> Decreasing chloride concentrations Stable <i>E. coli</i> and total suspended solid concentrations Increasing total phosphorus concentrations
EM001 Eagle Marsh Drain	Marginal ↔	Poor	<ul style="list-style-type: none"> Exceedances in chloride, <i>E. coli</i> (56%), total phosphorus (87%), total suspended solids and zinc (51%). Potential stressors include: quarry dewatering, agricultural and rural run-off 	<ul style="list-style-type: none"> Decreasing chloride and total suspended solids concentrations Stable <i>E. coli</i> concentrations Increasing total phosphorus concentrations
KD001 Krafts Drain	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (79%), total phosphorus (95%), total suspended solids and zinc Potential stressors include: rural and urban run-off Algae observed during summer months 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable chloride and <i>E. coli</i> concentrations Increasing total phosphorus concentrations
LB001 Low Banks Drain	Poor ↔	Poor	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i>, nitrate, total phosphorus (92%), total suspended solids and zinc Potential stressors include: agricultural and rural run-off Severe algae growth observed during summer months 	<ul style="list-style-type: none"> Stable chloride, <i>E. coli</i>, total phosphorus and total suspended solid concentrations

PA001 Point Abino Drain	Fair ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in copper, <i>E. coli</i> (55%), total phosphorus (82%), total suspended solids and zinc. Potential stressors include: agricultural and rural run-off Site is influenced by backflow from Lake Erie which is likely improving water quality 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable <i>E. coli</i> and total phosphorus concentrations Increasing chloride concentrations
SM001 Six Mile Creek	Poor ↓	Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (54%), total phosphorus (100%), total suspended solids and zinc Potential stressors include: agricultural and rural run-off 	<ul style="list-style-type: none"> Decreasing chloride and total suspended solids concentrations Stable <i>E. coli</i> concentrations Increasing total phosphorus concentrations
WD001 Wignell Drain	Poor ↔	Very Poor	<ul style="list-style-type: none"> Exceedances in chloride, copper, <i>E. coli</i> (64%), nickel, total phosphorus (100%), and total suspended solids and zinc Potential stressors include: quarry dewatering historic industrial pollution, agricultural and rural run-off Algae observed during summer months 	<ul style="list-style-type: none"> Decreasing total suspended solids concentrations Stable chloride and <i>E. coli</i> concentrations Increasing total phosphorus concentrations
WE000 Welland Canal	Good ↔	Insufficient Data	<ul style="list-style-type: none"> Exceedance total phosphorus Water source at this site is predominately from the Lake Erie 	<ul style="list-style-type: none"> Decreasing chloride concentrations Stable <i>E. coli</i>, total phosphorus and total suspended solid concentrations

4.7.2 LAKE ERIE TRIBUTARIES: HILSENHOFF BIOTIC INDEX RESULTS

HBI results ranged from *very poor* to *poor* at most Lake Erie tributary stations (**Table 8**). Results from biological assessments for these stations are illustrated in **Appendix J**. Sediment loading, reduced baseflow, lack of in-stream habitat, and nutrient enrichment are primary causes of impairment at these stations. Samples have not been collected from SM001 due to high water depth, channel morphology, and access restrictions.

4.7.3 LAKE ERIE TRIBUTARIES KEY FINDINGS

- Based on the 2018-2022 data, all the Lake Erie tributaries have total phosphorus exceedances. The most impacted of these tributaries is Wignell Drain which had median concentrations 10 times the Provincial Water Quality Objective. Wignell Drain watershed needs to be prioritized for Best Management Practice (BMPs) works to reduce phosphorus loads to Lake Erie.
- Nickel is not included in the WQI calculation; however, nickel concentrations were found to frequently exceed the PWQO at Beaver Dam Creek station BD001 and Wignell Drain station WD001. These nickel exceedances are likely from previous industrial landuse.

5.0 GROUNDWATER QUALITY MONITORING PROGRAM

5.1 NPCA GROUNDWATER MONITORING NETWORK

The NPCA Groundwater Monitoring Network (**Figure 11**) comprises of two components. The first component is the Provincial Groundwater Monitoring Network (PGMN) which is a partnership between the MECP and the Conservation Authorities of Ontario. The PGMN was initiated in 2001 and is a province-wide groundwater monitoring program designed to collect long-term baseline data on groundwater quantity and quality in special areas of interest. There are currently 470 ambient groundwater monitoring wells in the program. Groundwater is monitored through a network of 15 monitoring wells located throughout the NPCA watershed in locally significant hydrogeological areas. Most PGMN monitoring wells are sampled yearly for water geochemistry and all wells are instrumented with datalogging equipment which records hourly groundwater levels.

The second component of the NPCA Groundwater Monitoring Network is a network of 31 monitoring wells installed at 23 different locations across the NPCA watershed through a project between the NPCA and the Ontario Geological Survey (OGS). This project was partially funded through the Niagara Water Smart Grant program for the installation of monitoring wells in 2014 and 2015. Each of the 23 locations has a groundwater monitoring well installed at the top of bedrock in an aquifer zone commonly known as the Contact-Zone Aquifer. Three (3) of 23 locations have a set of nested monitoring wells installed at various depths within the overburden sediments. These wells were initially installed to investigate regional groundwater flow of five distinct features within the NPCA jurisdiction, which include:

- Three (3) buried bedrock valleys (Erigan Channel, Chippawa-Niagara Falls Channel and Crystal Beach Channel) and their groundwater relationships to Lake Erie, Lake Ontario and the Niagara River;
- The Fonthill-Kame Delta Complex/Twelve Mile Creek watershed area; and

- The Upper Welland River watershed.

After the completion of the OGS project in 2018, the NPCA has continued to sample water geochemistry yearly in 23 Contact-Zone Aquifer wells and all these wells are instrumented with datalogging equipment which records hourly groundwater levels to provide a better understanding of ambient groundwater conditions across the NPCA watershed.



Provincial Groundwater Monitoring Program Well W0000362-3 in Pelham

NPCA Groundwater Monitoring Network




Figure 11: Location of monitoring wells in the NPCA watershed

5.1.1 Groundwater Levels

The NPCA has been monitoring water levels at all PGMN wells since 2003 and other NPCA wells since 2014. Groundwater levels are typically at their highest during the late-winter and spring but begin to drop during the dry summer months and are lowest in the fall. There is also yearly variation in the groundwater levels which is dependent on precipitation. In dry years (such as 2016), water levels can drop substantially from their seasonal highs; and conversely the water level drops in wet years (2009) are not as substantial. PGMN monitoring wells each have water levels that are seasonally and yearly variable due to several factors (geologic formation the well is installed in, soils, precipitation, etc.). The results of this monitoring are found in **Appendix K**. The groundwater level data from the NPCA groundwater monitoring network will be used to help better understand the impacts of local/provincial scale drought events and its connection to climate change.

5.1.2 Groundwater Geochemistry

The NPCA Groundwater Monitoring Network is sampled yearly in the fall by the NPCA. Water chemistry sampling for the PGMN component of the NPCA Groundwater Monitoring Network began in 2006 where groundwater quality samples are collected by NPCA staff and laboratory analysis is provided by the MECP. The remaining NPCA wells have been sampled since 2014. From 2014 to 2018 these monitoring wells were sampled in partnership with the OGS to assist with a Southern Ontario ambient geochemistry program; the results of which are presented in the OGS Groundwater Resources Study 17 (Colgrove, L.M. and Hamilton, S. M., 2018). Starting in 2019, the 23 Contact-Zone Aquifer monitoring wells have been sampled for the same parameters as the PGMN monitoring wells and analyzed through private lab.

Groundwater quality samples are analyzed for general chemistry, nutrients, metals, and project specific parameters. Groundwater chemistry results are compared to the Ontario Drinking Water Standards (ODWS, MOE 2003) to provide an indication of overall groundwater quality across the NPCA. **Table 9** summarizes the health-related exceedances of the ODWS from 2018-2022 for the NPCA Groundwater Monitoring Network. All health related ODWS exceedances in PGMN wells are flagged by the MECP and are reported to the NPCA, Region of Niagara Public Health Department and local municipalities. Wells with reported exceedances are subsequently re-sampled by the MECP to confirm the initial exceedance. Based on the type and source of the exceedance these agencies formulate an action plan to protect human health. In the case of the non-PGMN wells, exceedances were reviewed through the OGS Ambient Geochemistry program (Colgrove, L.M. and Hamilton, S. M., 2018).

Table 9: NPCA Groundwater Monitoring Network stations (PGMN Wells and NPCA Wells) with Health-Related Exceedances of the ODWS (2018-2022). Blue text exceedances are caused by natural groundwater conditions and red text exceedances are caused by human influences.

Well ID Location	Well Type	Formation	Year				
			2018	2019	2020	2021	2022
W0000073 (Grimsby)	Bedrock	Guelph-Lockport	Sodium	Sodium	Sodium	Sodium	Sodium
W0000080 (West Lincoln)	Bedrock	Guelph-Lockport	Sodium Fluoride	Sodium Fluoride Uranium	Sodium Fluoride Uranium	Sodium Fluoride Uranium	Sodium Fluoride Uranium
W0000288 (Hamilton)	Bedrock	Guelph-Lockport	No Exceedance	No Exceedance	No Exceedance	No Exceedance	No Exceedance
Gainsborough (West Lincoln)	Bedrock	Guelph	No Exceedance	No Exceedance	No Exceedance	Fluoride	Sodium Fluoride
Welland (Welland)	Bedrock	Guelph	No Exceedance	No Exceedance	No Exceedance	No Exceedance	Sodium
Smith Road (Hamilton)	Bedrock	Guelph	No Exceedance	No Exceedance	No Exceedance	No Exceedance	Arsenic
Tyneside (Hamilton)	Bedrock	Guelph	No Exceedance	No Exceedance	No Exceedance	No Exceedance	Sodium
Oak Hall (Niagara Falls)	Bedrock	Guelph	Sodium	Sodium	Sodium	Sodium	Sodium
Glynn A Green (Pelham)	Bedrock	Lockport	Fluoride	Fluoride	Fluoride	Fluoride Arsenic	Sodium Fluoride Arsenic
W0000289 (Port Colborne)	Bedrock	Onondaga	No Exceedance	No Exceedance	No Exceedance	No Exceedance	No Exceedance

Well ID Location	Well Type	Formation	Year				
			2018	2019	2020	2021	2022
W0000341 (Lincoln)	Bedrock	Clinton	Sodium	Sodium	Sodium	Sodium	Sodium
W0000357 (Pelham)	Overburden	Fonthill Kame	No Exceedance	No Exceedance	Unable to sample	Unable to sample	Sodium
W0000361-2 (Pelham)	Overburden	Fonthill Kame	Nitrate	Nitrate	Sodium	Sodium Nitrate	Sodium Nitrate
W0000361-3 (Pelham)	Overburden	Fonthill Kame	Sodium	Sodium	Sodium	Sodium	Sodium
W0000362-2 (Pelham)	Overburden	Fonthill Kame	Sodium	Sodium	Sodium	Sodium	Sodium
W0000362-3 (Pelham)	Overburden	Fonthill Kame	Sodium	Sodium	Sodium	Sodium	Sodium
W0000384 (NOTL)	Overburden	Iroquois Sandplain	Sodium Nitrate	Sodium Nitrate	Sodium	Sodium	Sodium
Concession 1 (Wainfleet)	Bedrock	Bertie	Sodium	Sodium	Sodium	Sodium	Sodium
Stevensville (Fort Erie)	Bedrock	Bertie	No Exceedance	No Exceedance	No Exceedance	No Exceedance	Sodium
Monument (Haldimand County)	Bedrock	Bois Blanc	No Exceedance	No Exceedance	No Exceedance	No Exceedance	Sodium
North Pelham (Pelham)	Bedrock	Eramosa	N/A	No Exceedance	No Exceedance	No Exceedance	No Exceedance
5 th Street	Bedrock	Queenston	Sodium	No Exceedance	Sodium	Sodium	Sodium

Well ID Location	Well Type	Formation	Year				
			2018	2019	2020	2021	2022
(St.Catharines)							
Baden Powell (Niagara Falls)	Bedrock	Salina	Sodium	Sodium	Sodium	Sodium	Sodium
Townline (Port Colborne)	Bedrock	Salina	Sodium	Sodium	Sodium	Sodium	Sodium
Young Matthews (Niagara Falls)	Bedrock	Salina	No Exceedance	No Exceedance	No Exceedance	No Exceedance	Sodium
Buchner (Welland)	Bedrock	Salina	Sodium	Sodium	Sodium	Sodium	Sodium
College (Fort Erie)	Bedrock	Salina	Sodium Boron	Sodium Boron	Sodium Boron	Sodium Boron	Sodium Boron
Gents Road (Wainfleet)	Bedrock	Salina	No Exceedance	No Exceedance	No Exceedance	No Exceedance	Sodium
Sumbler Centre (Pelham)	Bedrock	Salina	No Exceedance	No Exceedance	No Exceedance	No Exceedance	Sodium
Public Works (Wainfleet)	Bedrock	Salina	Sodium	Sodium	Sodium	Sodium	Sodium
Road 9 (Haldimand County)	Bedrock	Salina	No Exceedance	No Exceedance	No Exceedance	No Exceedance	Sodium
Haldibrook Road (Haldimand County)	Bedrock	Salina	Sodium	Sodium	Sodium	Sodium	Sodium

Well ID Location	Well Type	Formation	Year				
			2018	2019	2020	2021	2022
Hart Road (Haldimand County)	Bedrock	Salina	Sodium	Sodium	Sodium	Sodium	Sodium
Case Bell (Wainfleet)	Bedrock	Salina	Sodium	Sodium	Sodium	Sodium	Sodium
W0000287 (Haldimand County)	Bedrock	Salina	Sodium	Sodium	Sodium	Sodium	Sodium
W0000290 (Niagara Falls)	Bedrock	Salina	Sodium Boron	Sodium Boron	Sodium Boron	Sodium Boron	Sodium Boron

5.1.3 KEY GROUNDWATER FINDINGS

- Exceedances of ODWS of arsenic (Glynn A Green & Smith Road), boron (W290-1 & College), fluoride (W080-1 & Glynn A Green) and uranium (W080-1) were observed in the NPCA sampling program, and these exceedances (except the Smith Road arsenic exceedance) have been attributed to natural groundwater conditions by the MECP Environmental Monitoring and Reporting Branch (EMRB) staff and by OGS Ambient Geochemistry program (Colgrove, L.M. and Hamilton, S. M., 2018). These elements occur naturally in the groundwater due to the dissolution of minerals from the bedrock formations. No anthropogenic activities or potential sites were identified. This appears to be an aquifer wide issue that will likely be present when this water is extracted for use. Regular water testing and treatment is recommended when using these aquifer sources for drinking. The new arsenic exceedance at the Smith Road well will be verified by additional NPCA confirmatory sampling.
- Most NPCA Groundwater Monitoring Network wells have sodium concentrations which exceed the 20 mg/L concentration. This is a health related ODWS exceedance where MECP has informed the Medical Officer of Health for the Health units so that this information could be communicated to local physicians for their use with patients on sodium restricted diets. MECP, EMRB and OGS Ambient Geochemistry program (Colgrove, L.M. and Hamilton, S. M., 2018) have attributed these exceedances to natural groundwater conditions and possibility from road salt in the shallow overburden wells. Regular water testing and treatment is recommended when using these aquifer sources for drinking.
- Elevated nitrate concentrations observed at monitoring wells W384-1 and W361-2 from 2003-2019 and 2021-2022 (W361-2 only) were attributed to agricultural landuse and/or faulty septic systems. In response to these exceedances, additional groundwater sampling of local private wells was completed by the NPCA in partnership with the Region of Niagara Public Health Unit in October 2008 and November 2009. The purpose of the additional sampling was to determine the extent of nitrate contamination near PGMN wells (W384-1 and W361-2) and to notify affected residents of potential health concerns related to elevated nitrate concentrations in drinking water. Sampling results indicated that none of the private wells tested exceeded the Ontario Drinking Water Standard (ODWS) for nitrate (MOE 2003) near W384-1 and one private well was found to exceed the ODWS near W361-2. The well exceeding the ODWS was determined to be a shallow dug well with poor construction and is likely not related to the nitrate exceedance at PGMN well W361-2. From 2018 to 2020 nitrate ODWS exceedances were not detected in both wells, and these reductions may be attributed to changes of local land uses and allowed for attenuation of nitrate. However, as 2021, nitrate concentrations have increased over Ontario Drinking Water Standards again at W361-2 and suggests a continue local source of nitrate.
- In general, ambient groundwater geochemistry across the NPCA was found to have elevated levels of chloride, iron, manganese sodium, and sulfate that exceed the Aesthetic Objectives within the ODWS (MOE 2003). The parameters which have Aesthetic Objectives may not be the source of any health-related issues, but these exceedances typically make the groundwater unpalatable or otherwise unpleasant for normal use (i.e., build-up/staining). Regular water

testing and treatment is recommended when using these aquifer sources for drinking and other uses.

5.2 WATER WELL DECOMMISSIONING PROGRAM

In 2022, the NPCA continued to provide grants to watershed residents interested in properly decommissioning abandoned water wells on their property through the NPCA Water Well Decommissioning Program. The grant program offers an 80% subsidy for water well decommissioning to a maximum of \$1000 per well. Grant applications are prioritized in areas designated as highly susceptible to groundwater contamination in the NPCA Groundwater Study (Waterloo Hydrogeologic Inc. 2005), areas where there is a high density of private wells used for domestic purposes, and areas where a watershed plan has been completed or is underway. Numerous improperly abandoned water wells are known to exist in the NPCA watershed, and these wells can serve as a direct pathway between potential contaminants at ground surface and deeper aquifers. The implementation of this program will reduce the risk of groundwater contamination and fulfills a recommendation made in the Groundwater Management Strategy of the NPCA Groundwater Study (Waterloo Hydrogeologic Inc. 2005). To date, 112 water wells have been decommissioned with the NPCA water well decommissioning program **Table 10**. An example of a water well decommissioning project is shown in **Figure 12**.

Table 10: Number and location of abandoned water wells decommissioned through the NPCA Water Well Decommissioning Grant from 2007 to 2022.

Year	# of Projects	Location of Projects
2007	4	Hamilton (2), Lincoln (1), Niagara Falls (1)
2008	1	Niagara-on-the-Lake (1)
2009	3	Grimsby (1), Lincoln (1), Niagara Falls (1)
2010	7	Grimsby (1), Lincoln (1), Pelham (3), St. Catharines (2), West Lincoln (1)
2011	9	Niagara Falls (1), NOTL (1), Pelham (2), Port Colborne (3), Wainfleet (1), West Lincoln (1)
2012	10	St. Catharines (1), NOTL (1), Pelham (1), Port Colborne (1), Wainfleet (1), West Lincoln (1), Fort Erie (2), Lincoln (2)
2013	12	St. Catharines (2), Niagara Falls (1), NOTL (3), Pelham (1), Lincoln (2), Wainfleet (2), West Lincoln (1)
2014	12	Niagara Falls (1), Fort Erie (1), NOTL (2), Pelham (3), Lincoln (1), Welland (2), Port Colborne (1), Thorold (1)
2015	9	NOTL (1), Pelham (3) Colborne (1), St. Catharines (2), Wainfleet (2)
2016	9	Hamilton (1), Lincoln (2), Niagara Falls (1), NOTL (1), Pelham (1), Wainfleet (3)
2017	8	Hamilton (2), Pelham (1), NOTL (1), Wainfleet (2), West Lincoln (2)
2018	10	Lincoln (4), NOTL (2), Pelham (1), St. Catharines (2), West Lincoln (1)
2019	5	Grimsby (1), Hamilton (1), Lincoln (1), Niagara Falls (1), Pelham (1)
2020	8	Fort Erie (1), Grimsby (1), Lincoln (1), NOTL (2), Pelham (1), Port Colborne (1), Thorold (1)
2021	5	Fort Erie (1), Hamilton (3), Pelham (1)
2022	5	Hamilton (2), St. Catharines (2), Welland (1)



Figure 12: An example of a NPCA Water Well Decommissioning Project. Left photo shows an abandoned dug well in need of decommissioning and the right photo shows same well after decommissioning had been completed by a licensed well contractor.

6.0 OTHER PROJECTS

6.1 HAMILTON INTERNATIONAL AIRPORT

Since 1998, the NPCA has been commissioned and funded by the Hamilton International Airport (HIA) to complete annual biological assessments of water quality near their property. The goal of the annual assessment is to determine if stormwater runoff and de-icing fluids such as propylene glycol are impacting surface water quality in two headwater tributaries of the Welland River. The NPCA generates this separate report for the HIA for their exclusive information and use.

6.2 GLANBROOK LANDFILL

Since 1998, the NPCA has been commissioned and funded by the City of Hamilton to complete biennial biological assessments of water quality for the Glanbrook Landfill. The Glanbrook Landfill is owned and operated by the City of Hamilton, and is designed to receive domestic, commercial, and non-hazardous solid industrial waste. The purpose of the biennial assessments is to determine if stormwater runoff and leachate from the landfill are negatively impacting water quality and aquatic biota in the Welland River and Buckhorn Creek. The NPCA generates this separate report for the City of Hamilton for their exclusive information and use.

6.3 UPPER TWELVE MILE CREEK TEMPERATURE MONITORING

The upper Twelve Mile Creek watershed contains the few identified coldwater streams in the NPCA watershed. Coldwater streams are fed by groundwater, and they are characterized by sustained cold water temperatures (approximately $<19^{\circ}\text{C}$) year-round. These streams often flow during dry periods of year because they are not dependent upon precipitation or other surface water. Water temperature is an important factor in determining the composition of aquatic communities. Coldwater sensitive species (e.g., Trout, Sculpins, some Mayflies and Stoneflies) can't breathe very efficiently and therefore need cold, clean water to survive. Therefore, these systems support a unique aquatic community adapted for coldwater conditions and significant warming can stress and even eliminate coldwater species from these watersheds. There are several threats to the health of coldwater streams and they include urbanization, climate change and overuse of groundwater resources. In 2013, the NPCA reinitiated temperature monitoring in the upper Twelve Mile Creek watershed to (1) identify and classify the thermal regime for the Twelve Mile Creek surface water sampling stations; and (2) identify any changes that may have occurred to the thermal stability of Twelve Mile Creek.

The NPCA monitored summer stream temperature at nine stream sites in 2022 (**Figure 13**). The stations were chosen due to the availability of background data, including water chemistry, benthic macroinvertebrate data, fisheries, stream morphology, hydrology data and stream temperature data. Onset HOBO Water Temp Pro dataloggers (**Figure 14**) were deployed at seven locations and stream temperature data was used from NPCA stream flow stations that are instrumented with temperature loggers. Hobo temperature loggers were deployed in June 2022 and collected in November 2022, recording at fifteen-minute intervals.

Stoneman and Jones (1996) developed a method to classify stream thermal stability with single observations of daily maximum air temperatures and water temperatures at 16:00 hours from June to September in given year. This method is by many Conservation Authorities in Ontario. The NPCA used June 4, 2022, to September 20, 2022, for the summer temperature monitoring period which is the warmest period during the year. To analyze the stream temperature data, the NPCA uses modified nomograms (Stoneman and Jones 1996) to determine stream thermal stability and estimate the thermal regime of each creek being monitored. This method determines whether a watercourse is to be classified as coldwater, coolwater or warmwater. Stoneman and Jones (1996) define the following 1) coldwater sites which support a large population of coldwater species; 2) coolwater sites that support a mix of coldwater-coolwater-warmwater; and 3) warmwater sites support few or no coldwater species.

Temperature monitoring locations




Figure 13: Map of temperature monitoring locations



Figure 14: Hobo Water Temperature Logger

Nomograms (**Figures 15 & 16**) were produced for the Effingham and St. John's tributary branches of the Upper Twelve Mile Creek from the NPCA 2022 stream temperature. Based on this analysis most (TW000, TW002, TW003, TW004, and Sulphur Springs) of the upper Twelve Mile Creek sites were classified as coldwater. TW001 TW005, TW006 and TW007 were classified as coolwater. No sites were classified as warmwater.




Figure 15: Thermal stability nomograms for 12 Mile Creek Effingham Tributary




Figure 16: Thermal stability nomograms for 12 Mile Creek St. John's Tributary and main channel

The classification changed at several sites (TW002, TW003, TW004, and Sulphur Springs) from coolwater to coldwater (**Table 11**) when compared to 2021 data. NPCA could not identify a specific reason for the change in classification but it likely a caused by yearly natural variation of stream temperatures. The classifications of these sites are consistently at the boundary of coldwater-coolwater interface and often transition between these classifications over the NPCA observations. Additional years of monitoring are required to determine if there are warming trends in these tributaries. The classifications of all other sites have generally remained stable with previous assessments.

Table 11: Stream classification summary 2011-2022. Black indicates no data, blue indicates coldwater, green indicates coolwater, and red indicates warmwater classifications.

Monitoring Site	2011	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
TW000	Cold Water	Cold Water	Cold Water	Cold Water	Cold Water	Cold Water	Cold Water	Cold Water	Cold Water	Cold Water	Cold Water
TW001	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water
TW002	Cool Water	Cool Water	Cold Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cold Water	Cool Water	Cold Water
TW003	Cool Water	Cold Water	Cold Water	Cold Water	Cool Water	Cool Water	Cool Water	Cold Water	Cold Water	Cool Water	Cold Water
Sulphur Springs					Cool Water	Cool Water	Cool Water	Cool Water	Cold Water	Cool Water	Cold Water
TW004	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cold Water
TW005	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water
TW006	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water	Cool Water
TW007	Cool Water	Warm Water			Warm Water	Cool Water		Warm Water	Warm Water	Warm Water	Cool Water

The NPCA will continue to monitor stream temperature as an additional method to evaluate the health and status of watershed communities in NPCA's significant coldwater watersheds.

6.4 LAKE NIAPENCO PERFLUORINATED COMPOUND MONITORING

Since 2012, the NPCA has been monitoring for Per- and polyfluoroalkyl substances (PFAS) in Lake Niapenco at Binbrook Conservation Area. PFAS were found in the plasma of snapping turtles at Lake Niapenco in 2009 and 2010 by an Environment Canada (EC) scientist as part of an organic toxins' accumulation study (de Solla et al. 2012). PFAS trackdown studies by MECP confirmed the presence of PFAS in Lake Niapenco and identified John C. Munro International Airport as the source of the contamination (Fowler 2011). Since 2015, Transport Canada and Procurement Canada have retained Arcadis Canada Inc. to conduct a risk assessment to investigate presence and distribution of PFAS in the Welland River downstream of the HIA.

NPCA collected a water sample at Lake Niapenco on June 20, 2022. **Figure 17** shows the sample location in the Conservation Area. The sampling protocol was as follows: 1) Lake Niapenco samples were collected in waist-deep water at the beach (**Figure 18**); and 2) Samples were collected and placed in a cooler with ice and shipped the next day for PFAS analysis.



Figure 17: Sample Location at Lake Niapenco



Figure 18: Sample location at the beach Lake Niapenco

Eurofins Lancaster Laboratories Environmental provided the results to the NPCA on June 20, 2022. The water chemistry results from Lake Niapenco indicated the present of PFOS at concentrations below Health Canada Provisional Drinking Water Guidelines (**Figure 19**). The concentration of PFOS generally matched the PFOS concentrations observed in previous NPCA sampling events but higher than the range of mean concentrations detected by de Solla *et al.* (2012) shown in **Figure 19**. Several other PFCs were detected in the Lake Niapenco. These included Perfluorooctanoic acid (PFOA), Perfluorohexane Sulfonate (PFHxS), Perfluorohexanoic Acid (PFHxA) and Perfluoropentanoic Acid (PFPeA) (**Figure 20**). These concentrations generally matched previous sampling event concentrations and PFPeA concentrations were higher than

those detected in the study by de Solla *et al.* (2012). The concentration of PFOA in Lake Niapenco was below Health Canada Provisional Drinking Water Guidelines. The other PFCs (PFHxS, PFHxA and PFPeA) have no drinking water guidelines.

Lake Niapenco was contaminated by historic PFAS usage from Hamilton International Airport (de Solla 2012 and Fowler 2011). The NPCA's monitoring at Binbrook Conservation Area continues to support the evidence that PFAS and specifically PFOS are present in Lake Niapenco but not at concentrations above Health Canada drinking water guidelines. It is expected that PFAS will continue to be present in Lake Niapenco due the persistence of PFAS in the environment and due to the delay to contain upstream sources. It should be noted that Hamilton Public Health has evaluated previous PFAS water quality data at Binbrook Conservation Area and determined that the PFAS concentrations detected would not adversely affect the park users.




Figure 19. PFOS concentrations found in Lake Niapenco by NPCA monitoring (2012-2022)




Figure 20: PFHxS, PFHxA, PFPeA and PFOA concentrations found in Lake Niapenco by NPCA (2012-2022)

6.5 CANADA ONTARIO AGREEMENT CLIMATE CHANGE MONITORING NETWORKS REVIEW PROJECT

A climate change sensitivity assessment completed by the MECP in 2009 identified the NPCA watershed as one of several southern Ontario watersheds that are highly vulnerable to the impacts of climate change. Indicators used in their analysis were related to water quality and quantity for both surface and groundwater resources. These indicators included frequency of low water levels, water use, water quality at active PWQMN stations, shallow well vulnerability, and baseflow. As a follow-up to this assessment, the NPCA conducted a detailed assessment in 2009-2010 of their existing monitoring networks and made specific recommendations for climate change detection and adaptation monitoring. Based on the NPCA and MECP assessments the existing monitoring station at Balls Falls Conservation Area was upgraded to an integrated monitoring site in 2015 which includes a rain gauge, soil moisture sensors and extended laboratory analysis for event sampling.

6.6 CONSERVATION AREA WATER QUALITY MONITORING

The NPCA Water Quality Monitoring Program was expanded in 2017 to include the waterbodies of the Conservation Areas: Dils Lake, Jordan Harbour, Lake Niapenco, Mud Lake, St. John's Pond, Virgil Reservoir, Wainfleet Wetlands and Wainfleet Bog. Water samples are collected quarterly or seasonally during the year and analyzed for general chemistry, nutrients, metals, and bacteria.

The water quality results for the NPCA Conservation Area locations are shown in **Figures 21-23**. Generally, the water quality observed in the Conservation Areas met environmental threshold values, but some exceedances were found. Total phosphorus exceeded the PWQO at most Conservation Areas, but Virgil Reservoir, Mud Lake and Wainfleet Bog had notably higher observed concentrations. The lentic Conservation Area environments (St. Johns, Binbrook, Mud Lake, Dils Lake and Virgil Reservoir) which have higher water temperatures, greater sunlight exposure, and reduced water flow are particularly vulnerable to excessive phosphorus concentrations which can stimulate the overgrowth and decomposition of plants and algae causing water quality concerns. There were chloride, copper, total suspended solids, and nitrate exceedances observed in Virgil Reservoir and these are likely due to the rural land uses, the proximity to Four Mile Creek Road and an abundance of carp in the reservoir. The pH of Wainfleet Bog also exceeded the lower threshold of the PWQO however this type of wetland is naturally acidic. This acidity is a by-product of microbial decay processes, cation exchange of the Sphagnum moss vegetation and input of additional acids from the atmosphere. Copper and zinc exceedances were observed at Wainfleet Bog main drainage ditch. This may be due past levels of atmospheric metal depositions that have accumulated in the peat soils and now being released in the surface water of the drainage ditch. The NPCA will continue to investigate the source of these metal exceedances.



Staff installing sampling equipment in Lake Niapenco




Figure 21: Box and Whisker Plot of chloride, total suspended solids, nitrate, and total phosphorus concentrations of the NPCA Conservation Area monitoring stations for 2018-2022 (n=14 for each station). The box length of the box-and-whisker plots represents the inter-quartile range that contains the median value shown as a horizontal line. The whiskers represent the minimum and maximum values. Outlier values represented by circles. Dashed lines represent Provincial Water Quality Objectives (PWQO), and Canadian Water Quality Guidelines (CWQG).







Figure 22: Box and Whisker Plot of pH, copper, lead, and zinc concentrations of the NPCA Conservation Area monitoring stations for 2018–2022 (n=14 for each station). The box length of the box-and-whisker plots represents the inter-quartile range that contains the median value shown as a horizontal line. The whiskers represent the minimum and maximum values. Outlier values represented by circles. Dashed lines represent Provincial Water Quality Objectives (PWQO).




Figure 23: Box and Whisker Plot of *E.coli* concentrations of the NPCA Conservation Area monitoring stations for 2018-2022 (n=14 for each station). The box length of the box-and-whisker plots represents the inter-quartile range that contains the median value shown as a horizontal line. The whiskers represent the minimum and maximum values. Outlier values represented by circles. Dashed lines represent Provincial Water Quality Objectives (PWQO).

6.7 LOWER TWELVE MILE CREEK TOTAL PCBs MONITORING

In 2020, MECP and City of St. Catharines had been investigating potential offsite impacts on surface water of Twelve Mile Creek from the former GM industrial area. To support agency partners, the NPCA has provided monthly samples (beginning September 2020) upstream and downstream of the former GM plant on 12 Mile Creek. This project included enhanced monitoring at the PWQMN station at Lakeport Road (TW009) and the reactivating a former monitoring site (TW011) at Welland Vale Road (See **Figure 24**). Samples collected are being analyzed for general chemistry, metals, nutrients, bacteria, and total PCBs.




Figure 24: Location of monitoring stations in upstream (TW011) and downstream (TW009) of former GM industrial area

Table 13: Twelve Mile Creek Surface water quality laboratory results (2022) from upstream (TW011) and downstream (TW009) of the former GM industrial Area

NPCA Site ID	Date	Alkalinity (mg/L)	pH	Solids; suspended (mg/L)	Chloride (mg/L)	E Coli (cfu/100mL)	Nitrate (mg/L)	Nitrite (mg/L)	phosphorus; total (mg/L)	Total PCBs (ug/L)	Aluminum (ug/L)	Barium (ug/L)	Beryllium (ug/L)	Bismuth (ug/L)	Cadmium (ug/L)	Chromium (ug/L)	Cobalt (ug/L)	Copper (ug/L)	Iron (ug/L)	Lead (ug/L)	Lithium (ug/L)	Manganese (ug/L)	Molybdenum (ug/L)	Nickel (ug/L)	Silver (ug/L)	Strontium (ug/L)	Tin (ug/L)	Titanium (ug/L)	Uranium (ug/L)	Vanadium (ug/L)	Zinc (ug/L)	Zirconium (ug/L)
PWQO CWQG			6.5-8.5	30	120	100	3	0.06	30	0	75		1100		0.2	1		5	300	5			40	25					5	6	30	4
Upstream GM TW011	2022-02-16	107	8.05	<3	23.8	260	0.19	<0.010	12.1	<0.04	39.2	22.1	<0.01	<0.005	<0.50	<0.10	0.94	50	0.06	2.3	3.29	1.05	0.78	<0.05	191	<0.10	0.75	0.39	<0.50	<3.0	<0.20	
Downstream GM TW009	2022-02-16	109	7.97	<3	27.9	34	0.22	<0.010	0.18	<0.04	62.8	22.4	<0.01	<0.005	<0.50	<0.10	0.99	80	0.14	2.2	5.9	1.06	0.84	<0.05	194	<0.10	1.23	0.4	<0.50	<3.0	<0.20	
Upstream GM TW011	2022-03-28	96.3	8.18	<3.0	21	310	0.28	<0.01	14	<0.04	94.1	19.1	<0.01	0.01	<0.50	<0.10	5.42	108	0.34	2	3.75	0.91	0.73	<0.05	174	<0.10	1.88	0.38	<0.50	6.4	<0.20	
Downstream GM TW009	2022-03-28	101	7.93	6.67	22.3		0.32	<0.01	22.9	<0.04	77.2	21.4	<0.1	<5	<0.9	<1	<1	1.07	83.2	<7	<5	6.5	<2	<2	<9	193	<9	1.61	3.79	<0.5	4.12	<1
Upstream GM TW011	2022-04-25	119	8.28		21.8	89	0.33	<0.01	12.8	<0.06	176	22.1	<0.02	0.01	<0.50	0.12	1.2	226	0.19	1.6	8.01	1.01	1.01	<0.01	203	<0.10	3.96	0.4	0.53	<3.0	<0.20	
Downstream GM TW009	2022-04-25	107	7.98	8.59	22.6	54	0.43	<0.01	16.4	<0.06	126	22.9	<0.1	<5	<0.9	<1	<1	1.04	99.4	<7	<5	8.02	<2	<2	<9	209	<9	3.82	<3	<0.5	4.4	<1
Upstream GM TW011	2022-05-30	97.5	8.17	10.5	18.2	140	0.2	<0.01	17.1	<0.06	234	21.5	<0.02	0.01	0.58	0.17	1.38	302	0.22	2.2	10.7	1.1	1.34	<0.01	176	<0.10	4.8	0.39	0.69	<3.0	<0.20	
Downstream GM TW009	2022-05-30	94.8	7.94	14.8	17.8	48	0.29	<0.01	23.3	<0.06	195	22.3	<0.1	<5	<0.9	<1	<1	1.16	164	<7	<5	10.3	<2	<2	<9	177	<9	6.99	<3	<0.5	7.09	<1
Upstream GM TW011	2022-06-27	95.1	8.34	9.4	17.8		0.12	<0.01	16.3	<0.06	262	21.8	<0.02	0.01	<0.50	0.17	1.22	329	0.25	1.8	11.2	1.04	1.05	<0.01	163	<0.10	5.66	0.36	0.68	<3.0	<0.20	
Downstream GM TW009	2022-06-27	94	7.94	15.7	16.9	480	0.17	<0.01	24.8	<0.06	311	23.9	<0.1	<5	<0.9	<1	<1	1.01	207	<7	<5	12.6	<2	<2	<9	170	<9	10.9	3.57	0.56	7.43	<1
Downstream GM TW009	2022-07-25	97.1	8	11.4	18.4		0.16	<0.01	22.2	<0.06	132	23.5	<0.1	<5	<0.9	<1	<1	1.02	119	<7	<5	10.2	<2	<2	<9	175	<9	4.28	3.25	<0.5	5.63	<1
Upstream GM TW011	2022-08-18	99.5	8.27	4.2	18.8	420	0.07	<0.01	13.9	<0.06	142	22.3	<0.02	0.01	0.5	0.11	1.03	179	0.15	1.7	7.18	1.16	0.95	<0.01	176	<0.10	3.1	0.38	0.59	<3.0	<0.20	
Downstream GM TW009	2022-08-18	103	8.25	3.6	18.5	200	0.07	<0.01	15.2	<0.06	130	22.2	<0.02	5	<0.50	<0.10	1.02	168	0.15	1.7	8.9	1.14	0.87	<0.01	174	<0.10	<3.00	0.38	0.56	<3.0	<0.20	
Upstream GM TW011	2022-10-24	101	8.26	7.6	18.4	>200	0.12	<0.01	18.6	<0.06	163	22	<0.02	0.01	<0.50	0.12	0.98	230	0.18	1.6	8.7	1.12	1.05	<0.01	179	<0.10	3.68	0.38	<0.50	<3.0	<0.20	
Downstream GM TW009	2022-10-24	101	7.99	8.02	17.6	180	0.15	<0.01	18.9	<0.06	101	23.2	<0.1	<5	<0.9	<1	<1	<0.5	88.4	<7	<5	6.94	<2	<2	<9	181	<9	3.25	3.47	<0.5	4.22	<1
Upstream GM TW011	2022-11-28	104	8.42	6.1	18.2	129	0.1	<0.01	13.2	<0.06	156	22.3	<0.02	0.01	<0.50	0.1	0.98	196	0.17	1.7	7.15	1.1	0.87	<0.01	192	0.19	<6.00	0.37	<0.50	<3.0	<0.20	
Downstream GM TW009	2022-11-28	102	7.9	9.53	19.5	163	0.23	<0.01	19.7	<0.06	62.2	19.9	<0.1	<5	<0.9	<1	<1	1.54	94.9	<7	<5	8.61	<2	3.15	<9	186	<9	1.68	>3	<0.5	4.42	<1
Upstream GM TW011	2022-12-13	102	8.32	5.9	17.6	400	0.13	<0.01	14	<0.06	178	23.2	<0.02	0.01	0.72	0.11	1.03	219	0.17	1.8	7.11	1.12	0.99	<0.01	186	<0.10	4.14	0.39	0.61	<3.0	<0.20	
Downstream GM TW009	2022-12-13	102	8.05	6.59	17.8	98	0.18	<0.01	16.4	<0.06	159	21.6	<0.1	<5	<0.9	<1	<1	1.08	110	<7	<5	5.72	<2	<2	<9	176	<9	5.71	<3	<0.5	5.05	<1

Generally, in 2022, the water quality observed met environmental threshold values, but some exceedances were found. *E. coli* concentrations were found to routinely exceed the PWQO at both stations and sources possibly could include wildlife or human sources. There were some exceedances of metals observed, which are from a combination of natural and anthropogenic sources. Aluminum routinely exceeded PWQO and this likely due to the contribution of naturally elevated levels aluminum found in soils of the watershed. Iron (May & June 2022) and a copper (March 2022) exceedances at TW011 were observed, but these exceedances were not higher than concentrations historically observed by the NPCA in the upper and less-impacted portions of the Twelve Mile Creek watershed. All total PCBs results for both sites were less than laboratory detection in 2022 (**Table 13**)

Based on the NPCA data there is not sufficient information to conclude that runoff from the former GM industrial area is impacting Twelve Mile Creek at Lakeport Road. The NPCA will continue to conduct water quality sampling at these stations and provide data to support the investigation of potential offsite impacts of surface water of Twelve Mile Creek from the former GM industrial area.

6.8 NPCA DATA REQUESTS

The NPCA Water Quality Monitoring Program generates a large wealth of scientific data that is a valuable resource to several clients. In 2022 the NPCA water quality monitoring program received 45 data requests from a variety of agencies and the public. These include:

- Ontario Ministry of the Environment Conservation and Parks
- Ontario Ministry of Agriculture, Food and Rural Affairs
- Ontario Ministry of Natural Resources
- Academia (McMaster University & University of Waterloo)
- Environment Canada
- Municipalities (Upper and Lower Tier)
- Health Units (Hamilton and Niagara)
- Consultants
- Non-Governmental Agencies
- Public



NPCA staff collecting a grab sample for laboratory analysis.

7.0 CONCLUSIONS

Based on the foregoing, the NPCA offers the following conclusions:

- Based on the results of the 2018 to 2022 WQI, 59% of the NPCA surface water monitoring stations are rated as *poor*, 31% are rated as *marginal*, 9% are rated as *fair*, and 1% is rated as *good*.
- Based on the results of the 2018 to 2022 biological assessments using the Hilsenhoff Biotic Index (HBI): 23% of the NPCA monitoring stations had water quality rated as *very poor*, 38% rated as *poor*, 20% rated as *fairly poor*, 3% rated as *fair* and 16% have not been assessed.
- Generally, the WQI ratings at water quality stations were relatively stable when compared to historic NPCA data. The Welland Canal (Port Colborne) has the highest water quality rating in the NPCA watershed. This is not unexpected as the water found in the Welland Canal is from Lake Erie. The other monitoring sites with high WQI ratings include the Point Abino Drain (Fort Erie), the Effingham tributary of Twelve Mile Creek (Pelham), the upper Welland River (Hamilton), Beaver Dams Creek (Thorold) and Beaver Creek (Fort Erie).
- The NPCA watershed has total phosphorous exceedances of the Provincial Water Quality Objective at virtually all monitoring stations owing to the higher population densities, and larger concentration of agriculture and industry. Based on the data collected to date, elevated concentrations of total phosphorus are the most frequent (over 95% observations) and widespread cause of water quality impairment in the NPCA watershed. The relative high frequency and magnitude of these exceedances is the driving factor in lowering the WQI at all stations.
- Exceedances of *E. coli* also contribute greatly to lower WQI ratings in the NPCA watershed. Approximately 60% of the NPCA stations have median *E. coli* concentrations greater than the PWQO. *E. coli* concentrations in the 5 watersheds (One Mile Creek, Two Mile Creek, Walkers Creek and Prudhommes Drain) are high relative to other watersheds and the sources of these exceedances need to be examined further. The NPCA initiated a trackdown of *E. coli* sources in Two Mile Creek Conservation Area and discovered a storm sewer outfall as the likely source of the *E. coli* contamination. This information has been provided to the Town of the Niagara-on-the-Lake and the Town staff have implemented infrastructure upgrades to eliminate *E. coli* sources. The NPCA will continue to work with municipalities to identify *E. coli* sources.
- WQI ratings and Hilsenhoff Biotic Index results did not agree at every station (i.e., where the WQI rating is *marginal* the Hilsenhoff Biotic Index rating is *very poor*) indicating that the benthic invertebrate data does not entirely support the chemical data. There may be other factors which are beyond the scope of this analysis such as the availability of in-stream habitat, size of the dataset used to calculate the WQI rating, and influence of parameters not monitored by the NPCA that may be affecting this agreement. Nonetheless biological and chemical monitoring remain important tools to evaluating water quality.
- Exceedances for chloride, metals (copper, lead, and zinc), nitrate, and total suspended solids were uncommon in the NPCA watershed. Elevated copper exceedances in Richardsons Creek (St. Catharines) and Eighteen Mile Creek (Lincoln) warrants further investigation. Zinc exceedances in the Welland River are related to Hamilton Airport operations and the NPCA and MECP are working with the HIA to reduce concentrations. Chloride exceedances are related to road

salt impacts and groundwater discharge to surface water. Nitrate exceedances in the Richardson Creek and the upper Twelve Mile Creek may be related nearby commercial operations in the area. Total suspended solids are mainly related wet weather events and inadequate riparian buffers along watercourses.

- Most NPCA groundwater monitoring wells exceeded 20 mg/L ODWS for sodium which is a health-related concern for individuals with sodium restricted diet intakes. A small number of wells were found to have ODWS exceedances in arsenic, boron, fluoride, nitrate, and uranium. In addition, ambient groundwater geochemistry across the NPCA was found to have elevated levels of sulfate, iron, manganese, and chloride that exceed the Aesthetic ODWS. ODWS exceedances have been investigated by MECP and Public Health Units and are attributed to natural conditions of the groundwater or in the case of nitrate a local source. It is recommended that watershed residents using groundwater test their water regularly and treat water appropriately.
- The water temperature monitoring of the Upper Twelve Mile Creek headwaters classified five stations as coldwater and three stations as coolwater. The Upper Twelve Mile Creek watershed is capable of supporting coldwater species based on temperatures monitoring data however there is warming of the main channel of Twelve Mile as it travels through Short Hills Provincial Park where stream temperature transitions to a warm water classification.
- The NPCA Water Quality Monitoring Program continues to generate a large wealth of scientific data that is a valuable resource to the public, environmental consultants, community groups, educational institutions, and other governmental agencies. In addition, this program continues to provide technical support to other NPCA programs, including Technical, Stewardship and Development Services.

8.0 RECOMMENDATIONS

Recommendations from this NPCA Water Quality Monitoring Program Report are summarized as follows:

1. It is recommended that the NPCA continue the Water Quality Monitoring Program to collect up-to-date and reliable water quality data and continue to make this information freely accessible to the public.
2. It is recommended that the NPCA continue to analyze all collected water quality data with the intent to identify significant trends or abnormalities.
3. It is recommended that the NPCA continue to work with our partner municipalities and the MECP to identify and mitigate abnormally high sources of water pollution as they are identified through the Water Quality Monitoring field sampling program.
4. It is recommended that the NPCA continue to monitor summer water temperatures for the upper Twelve Mile Creek sampling stations to classify their thermal regimes and to identify any changes to the thermal stability of this watershed.

5. It is recommended that the NPCA continue to undertake annual water quality assessments for the Hamilton International Airport and the City of Hamilton's Glanbrook Landfill (both presently commissioned and funded by the City of Hamilton).
6. It is recommended that the NPCA continue to offer the 'Water Well Decommissioning Program' to the public to help reduce the risk of groundwater contamination by removing old and abandoned wells.
7. As it is no longer typical to have watercourses completely frozen from December to March, it is recommended that the NPCA continue to pursue opportunities to expand the surface water quality monitoring program outside of the months of April to November to address water quality data gaps which presently exist for the winter months.
8. It is recommended that the NPCA continue to monitor Per- and polyfluoroalkyl substances in Lake Niapenco to provide the public with annual reporting information. Yearly surveillance is an appropriate time interval for monitoring based on lab analysis costs and contaminant variability and concentrations. It is also recommended that NPCA continue to partner and share information with Transport Canada and the Risk Assessment Team to fully understand the presence and distribution of PFAS in the Welland River downstream of the HIA.
9. It is recommended that the NPCA continue to support and MECP Climate Change project for Twenty Mile Creek at Balls Falls Conservation Area by continuing to operate and maintain this enhanced monitoring station.

9.0 REFERENCES CITED

Canadian Council of Ministers of the Environment. 2001. Canadian Water Quality Index.

Canadian Council of Ministers of the Environment (CCME). 2002. Canadian water quality guidelines for the protection of aquatic life: Total particulate matter. In: Canadian Environmental Quality Guidelines, Canadian Council of Ministers of the Environment, Winnipeg

Canadian Council of Minister of the Environment. 2003. Canadian Water Quality Guidelines for the Protection of Aquatic Life. Canadian Environmental Quality Guidelines.

Canadian Council of Ministers of the Environment. 2011. Canadian Water Quality Guidelines for the Protection of Aquatic Life. Canadian Environmental Quality Guidelines.

Colgrove, L.M. and Hamilton, S.M. 2018. Geospatial distribution of selected chemical, bacteriological and gas parameters related to groundwater in southern Ontario; Ontario Geological Survey, Groundwater Resources Study 17.

Conservation Ontario. 2011. Guide to Developing Conservation Authority Watershed Report Cards. Newmarket.

de Solla, S., A. De Silva and R Letcher. 2012. Highly elevated levels of perfluorooctane sulfonate and other perfluorinated acids found in biota and surface water downstream of an international airport, Hamilton, Ontario, Canada. *Environment International* 39:19-26.

- Fowler, C. 2011. PFOS in the Welland River and Lake Niapenco. Ontario Ministry of the Environment. Technical Support Section, West Central Region Hamilton.
- Griffiths, R.W. 1999. Hilsenhoff Biotic Index: Bioassessment of Water Quality. The Centre for Environmental Training, Niagara College: Niagara-on-the-Lake, Ontario.
- Health Canada. 2016. Perfluorooctanoic Acid (PFOA) in Drinking Water.
<https://www.canada.ca/content/dam/hc-sc/healthy-canadians/migration/health-system-systeme-sante/consultations/acide-perfluorooctanoic-acid/alt/perfluorooctanoic-eng.pdf>
- Health Canada. 2016. Perfluorooctane Sulfonate (PFOS) in Drinking Water.
<https://www.canada.ca/content/dam/hc-sc/healthy-canadians/migration/health-system-systeme-sante/consultations/perfluorooctane-sulfonate/alt/perfluorooctane-sulfonate-eng.pdf>
- Helsel, D., and R. Hirsch. 1992. Statistical Methods in Water Resources, Elsevier, 522 p.
- Helsel, D., D. Muller, and J. Slack. 2005. Computer Program for the Kendall Family of Trend Tests, Scientific Investigation Report, USGS, 1-4.
- Hilsenhoff, William L. 1987. An Improved Biotic Index of Organic Stream Pollution. The Great Lakes Entomologist. 20: 31-36.
- Jones, C., K. Somers, B. Craig, and T. B. Reynoldson. 2005. Ontario Benthos Biomonitoring Network Protocol Manual: Version 1.0. Ontario Ministry of Environment: Dorset, Ontario.
- Jones, C. 2006. Personal communication. Benthic Biomonitoring Scientist, Environmental Monitoring and Reporting Branch, Ontario Ministry of the Environment.
- MacRitchie, S. 2009. Fluoride and Uranium Occurrence in PGMN Well 80-1. Ministry of the Environment: Environmental Monitoring and Reporting Branch.
- Ontario Ministry of the Environment and Environment Canada. 1995. Niagara River Remedial Action Plan Stage 2 Report.
- Ontario Ministry of the Environment. 1994. Water Management, Policies, Guidelines and Provincial Water Quality Objectives. Government of Ontario, Toronto.
- Ontario Ministry of the Environment. 2003. Technical Support Document for Ontario Drinking Water Standards, Objectives and Guidelines. Government of Ontario, Toronto.
- Regional Municipality of Niagara. 2003. Niagara Water Quality Protection Strategy. Prepared for the Regional Municipality of Niagara by MacViro, Philips Engineering and CH2MHill.
- Stoneman, Christine, and Michael Jones. 1996. A Simple Method to Classify Stream Thermal Stability with Single Observations of Daily Maximum Water and Air Temperatures. North American Journal of Fisheries Management. 16: 728-737.
- Tchobanoglous, G, and E. Schroeder. 1987. Water Quality. Addison-Wesley Publishing Company: Reading, Massachusetts.
- Waterloo Hydrogeologic Inc., 2005. NPCA Groundwater Study. Waterloo, Ontario.


10.0 ACKNOWLEDGEMENTS

The NPCA would like to thank the City of Hamilton, Ontario Ministry of Environment, Conservation and Parks, Regional Municipality of Niagara, and Haldimand County for their support of the NPCA Water Quality Monitoring Program.

Appendix A

Niagara Peninsula Conservation Authority

Water Quality Ratings 2018 - 2022



Surface Water Quality Stations
- WQI

Waterbodies
NPCA Jurisdiction
Sampled_Subwatersheds


Fair
Good
Marginal
Poor

0 5 10 20 30
Scale: 1:230,000
Kilometers


All Frames: North American Datum 1983, Universal Transverse Mercator 6° Projection, Zone 17N, Central Meridian 81° West.
Produced by the Niagara Peninsula Conservation Authority with data supplied under license by members of the Ontario Geospatial Data Exchange, 2017.




Appendix B




Welland River Watershed Chloride Concentrations 2018–2022




Welland River Tributaries Chloride Concentrations 2018–2022




Twenty Mile Creek Watershed Chloride Concentrations 2018–2022




Lake Ontario South Shore Tributaries Chloride Concentrations 2018–2022



Niagara River Tributaries Chloride Concentrations 2018–2022




Lake Erie North Shore Tributaries Chloride Concentrations 2018–2022



Appendix C

Niagara Peninsula Conservation Authority

Median Copper Concentrations 2018 - 2022



Surface Water Quality Stations
- Copper (ug/L)

- < 2.5
- 2.5 - 5.0
- 5.1 - 10

■ Waterbodies

■ NPCA Jurisdiction

■ Sampled Subwatersheds

Scale: 1:230,000


0 3.75 7.5 15 22.5 30

Kilometers


All Frames: North American Datum 1983, Universal Transverse Mercator 6° Projection, Zone 17N, Central Meridian 81° West.
Produced by the Niagara Peninsula Conservation Authority with data supplied under licence by members of the Ontario Geospatial Data Exchange, 2017.




Lake Ontario South Shore Tributaries Copper Concentrations 2018–2022




Lake Erie North Shore Tributaries Copper Concentrations 2017–2021




Lake Erie North Shore Tributaries Copper Concentrations 2018–2022




Twenty Mile Creek Watershed Copper Concentrations 2018–2022



Welland River Watershed Copper Concentrations 2018–2022




Welland River Tributaries Copper Concentrations 2018–2022



Appendix D

Niagara Peninsula Conservation Authority

Median E.coli Concentrations 2018 - 2022



Surface Water Quality Stations - E.coli (cfu/100mL)

- < 25
- 25 - 100
- 101 - 500
- > 500


- Watercourses
- Waterbodies
- Sampled Subwatersheds
- NPCA Jurisdiction

0 5 10 20 30
Scale: 1:230,000
Kilometers


All Frames: North American Datum 1983, Universal Transverse Mercator 6° Projection, Zone 17N, Central Meridian 81° West.
Produced by the Niagara Peninsula Conservation Authority with data supplied under license by members of the Ontario Geospatial Data Exchange, 2017.




Lake Ontario South Shore Tributaries E.coli Concentrations 2018-2022




Lake Erie North Shore Tributaries E.coli Concentrations 2018–2022




Niagara River Tributaries E.coli Concentrations 2018–2022




Twenty Mile Creek Watershed E.coli Concentrations 2018–2022




Welland River Watershed E.coli Concentrations 2018–2022




Welland River Tributaries E.coli Concentrations 2018–2022




Appendix E




Lake Erie North Shore Tributaries Lead Concentrations 2018–2022




Niagara River Tributaries Lead Concentrations 2018–2022




Lake Ontario South Shore Tributaries Lead Concentrations 2018–2022




Twenty Mile Creek Watershed Lead Concentrations 2018–2022



Welland River Tributaries Lead Concentrations 2018-2022




Welland River Watershed Lead Concentrations 2018–2022



Appendix F

Niagara Peninsula Conservation Authority

Median Nitrate Concentrations 2018 - 2022




Surface Water Quality Stations
- Nitrate ($\mu\text{g/L}$)

Waterbodies
Sampled Subwatersheds
NPCA Jurisdiction


Green: < 500
Yellow: $500 - 1300$
Red: $1301 - 2900$
Black: > 2900

0 5 10 20 30
Scale: 1:230,000
Kilometers


All Frames: North American Datum 1983, Universal Transverse Mercator 6° Projection, Zone 17N, Central Meridian 81° West.
Produced by the Niagara Peninsula Conservation Authority with data supplied under license by members of the Ontario Geospatial Data Exchange, 2017.




Lake Erie North Shore Tributaries Nitrate Concentrations 2018–2022




Lake Ontario South Shore Tributaries Nitrate Concentrations 2018–2022




Twenty Mile Creek Watershed Lead Concentrations 2018–2022




Niagara River Tributaries Lead Concentrations 2018–2022



Welland River Tributaries Nitrate Concentrations 2018–2022




Welland River Watershed Nitrate Concentrations 2018–2022



Appendix G

Niagara Peninsula Conservation Authority

Median Total Phosphorus Concentrations 2018 - 2022



Surface Water Quality Stations - Total Phosphorus (ug/L)


Waterbodies
Sampled Subwatersheds
NPCA Jurisdiction

0 5 10 20 30
Scale: 1:230,000
Kilometers


All Frames: North American Datum 1983, Universal Transverse Mercator 6° Projection, Zone 17N, Central Meridian 81° West.
Produced by the Niagara Peninsula Conservation Authority with data supplied under license by members of the Ontario Geospatial Data Exchange, 2017.




Lake Erie North Shore Tributaries Total Phosphorus Concentrations 2018–2022




Lake Ontario South Shore Tributaries Total Phosphorus Concentrations 2018–2022




Welland River Tributaries Total Phosphorus Concentrations 2018–2022




Twenty Mile Creek Watershed Total Phosphorus Concentrations 2018–2022



Welland River Tributaries Total Phosphorus Concentrations 2018-2022




Niagara River Tributaries Total Phosphorus Concentrations 2018–2022



Appendix H

Niagara Peninsula Conservation Authority


Median Total Suspend Solids Concentrations 2018-2022



Surface Water Quality Stations
- TSS (mg/L)

- < 10
- 10 - 20
- 21 - 40
- > 40


- Waterbodies
- Sampled Subwatersheds
- NPCA Jurisdiction




All Frames: North American Datum 1983, Universal Transverse Mercator 6° Projection, Zone 17N, Central Meridian 81° West.
Produced by the Niagara Peninsula Conservation Authority with data supplied under license by members of the Ontario Geospatial Data Exchange, 2017.




Lake Erie North Shore Tributaries Total Suspended Solids Concentrations 2018–2022




Lake Ontario South Shore Tributaries Total Suspended Solids Concentrations 2018–2022




Welland River Tributaries Total Suspended Solids Concentrations 2018–2022




Twenty Mile Creek Watershed Total Suspended Solids Concentrations 2018–2022




Welland River Watershed Total Suspended Solids Concentrations 2018–2022




Niagara River Tributaries Total Suspended Solids Concentrations 2018–2022




Appendix I




Lake Erie North Shore Tributaries Zinc Concentrations 2018–2022




Lake Ontario South Shore Tributaries Zinc Concentrations 2018–2022




Welland River Tributaries Zinc Concentrations 2018–2022




Twenty Mile Creek Watershed Zinc Concentrations 2018–2022



Welland River Watershed Zinc Concentrations 2018–2022



Niagara River Tributaries Zinc Concentrations 2018–2022



Appendix J




-  Fair
-  Fairly Poor
-  Poor
-  Very Poor
-  Insufficient

All Frames: North American Datum 1983, Universal Transverse Mercator 6° Projection, Zone 17N, Central Meridian 81° West.
Produced by the Niagara Peninsula Conservation Authority with data supplied under license by members of the Ontario
Geospatial Data Exchange, 2017.




Appendix K


Provincial Groundwater Monitoring Network Well W0000073-1




Provincial Groundwater Monitoring Network Well W0000080-1




Provincial Groundwater Monitoring Network Well W0000287-1




Provincial Groundwater Monitoring Network Well W0000288-1




Provincial Groundwater Monitoring Network Well W0000289-1



Provincial Groundwater Monitoring Network Well W0000290-1




Provincial Groundwater Monitoring Network Well W0000341-1




Provincial Groundwater Monitoring Network Wells W0000356-2 and W0000356-3


Corrected and Adjusted PGMIS Water Level Elevation (masl)




Provincial Groundwater Monitoring Network WellW0000357-1




Provincial Groundwater Monitoring Network Well W0000361-2




Provincial Groundwater Monitoring Network Well W0000361-3




Provincial Groundwater Monitoring Network Well W0000362-2




Provincial Groundwater Monitoring Network Well W0000362-3




Provincial Groundwater Monitoring Network Well W0000384-1



NPCA 5th Street Monitoring Well



NPCA BadenPowell Monitoring Well




— Levellogger Water Level

● Manual Measurements


NPCA Buchner Monitoring Well



NPCA CaseBell Monitoring Well



NPCA College Monitoring Well




— Levellogger Water Level

● Manual Water Level

NPCA Concession1 Monitoring Well




NPCA Gainsborough Monitoring Well




— Levellogger Water Levels

● Manual Water Levels

NPCA Gents Monitoring Well




NPCA Glynn A Green 20ft Monitoring Well




NPCA Glynn A Green 78ft Monitoring Well



NPCA Glynn A Green 112ft Monitoring Well




NPCA Glynn A Green 135ft Monitoring Well




NPCA Glynn A Green Monitoring Well




NPCA Haldibrook Monitoring Well




NPCA Hart Monitoring Well




NPCA Monument Monitoring Well




NPCA North Pelham Park Monitoring Well



NPCA Oak Hall Monitoring Well




NPCA Public Works Monitoring Well




— Levellogger Water Levels

● Manual Water Levels


NPCA Road9 Monitoring Well




NPCA Smith Monitoring Well




NPCA Smith 21ft Monitoring Well




NPCA Smith 60ft Monitoring Well



NPCA Stevensville Monitoring Well




NPCA SumblerCentre Monitoring Well




— Levellogger Water Level ● Manual Water Level


NPCA Townline Monitoring Well



NPCA Townline 25ft Monitoring Well




NPCA Townline 65ft Monitoring Well




Levellogger Water Levels

Manual Water Levels

NPCA Tyneside Monitoring Well



NPCA Welland Monitoring Well



NPCA YoungMatthews Monitoring Well

